

## DRAGONFLY GARDENS

www.dragonflygardens.net 715-554-7219 natives@dragonflygardens.net Native Catalog 2021 Edition 491 Hwy 46 Amery, WI

AVAILABILITY, SIZE and PRICE subject to change.

	Τ	1	I		AVAILABILITY, SIZE and PRICE subject to	2.101	50.	+
	Flowers	Ht.	Moisture	l iaht	Description	2" Pot	4" Pot	6" Pot 4
Wildflowers		J111C.	INIOISTUIE	Ligit	Description	7	4	ၑ
			_					
Achillea millefolium Yarrow	White June to Oct	18-30"	D, MD, M, MW	full sun to part shade	Attractive, aromatic, ferny foliage. Great component of a no- mow or low-mow lawn. Good butterfly plant. Grows well in a wide variety of habitats.	•		
Acorus calamus Sweet Flag	Yellow July to Aug	36"	MW, W	full sun to part shade	Unusual plant with attractive yellowish foliage and sweet fragrance when crushed. Forms beautiful clumps on lake margins. Great for water gardens.	•		
Agastache foeniculum Anise Hyssop	Blue June to Oct	24-40"	MD, M	full sun to part shade	Beautiful 3" flower spikes attract butterflies throughout summer. Excellent cutflower and nice in perennial gardens. Fragrant foliage makes good herbal tea.	•	•	
Allium cernuum Nodding Wild Onion	Pink July to Sept	12-18"	MD, M, MW	full to part sun	Nodding 2" flower clusters above thin, wispy stems. Edible flower bulbs.	•	•	
Allium stellatum Prairie Onion	Pink July to Sept	12-18"	D, MD, M, MW	full to part sun	Attractive plants occurring in dry prairies and on rocky cliff areas. Good plant for tough sites and rock gardens. Ranging throughout MN and WI.	•		
Allium tricoccum Wild Leek	White June to July	8"	MD, M, MW	full to part shade	Also known as "Ramps", edible leaves and bulbs are popular among culinary artists. They are very slow to reproduce so be patient and don't overharvest.		3"	
Amorpha canescens Leadplant	Blue June to July	18-42"	D, MD, M	full sun to part shade	Long-lived shrub with attractive fine silvery foliage. Slow- growing but worth the wait. Used medicinally by Native Americans. Suited to driest sandy soil.	•	•	
Anaphilis margaritacea Pearly Everlasting	White June to Aug	18-24"	D, MD	full sun to part shade	White papery strawflowers produced atop attractive whitish foliage. Nice dried flower and contrast plant in the perennial garden. Host plant for painted lady butterfly.	•	•	
Anemone canadensis Canada Anemone	White May to July	12-24"	M, MW	full sun to part shade	Fine, showy clump-forming plant of mesic to moist prairies, forest margins and woodlands.	•		
Anemone cylindrica Thimbleweed	White Late May to July	16-20"	D, MD, M	full sun to part shade	Thimble-like infloresence held atop slender stems above highly divided basal leaves. Very pretty from summer into fall. Good on very dry sandy soils.	•	•	
Anemone patens Pasque Flower	Light Blue April to May	6"	D, MD	full to part sun	One of the first wildflowers in spring. Very showy 3" flowers followed by a cluster of hairy fruits. Dry soils. Great in rock gardens.	•		
Aquilegia canadensis Wild Columbine	Orange May to June	20-36"	D, MD, M	full sun to shade	Free-flowering woodland perennial that forms large clumps with attractive foliage. Excellent hummingbird plant. Often referred to as "honeysuckle."	•	•	
Arisaema triphyllum Jack in the Pulpit	Purple/ Green May to June		MD, M, MW	part sun to shade	Very unusual flower stalk that emerges separately from 3- parted leaf. Great species for moist to wet forest restorations. Reliably spreads from seed once established.		3"	
Artemesia ludoviciana Prairie Sage	Yellow July to Oct	18-30"	D, MD, M	full to part sun	Fragrant whitish foliage makes a nice contrast to green plants. Dry to mesic prairies. Sacred incense and insect repellant of Native Americans.	•		
Asarum canadense Wild Ginger	Maroon May to June	6-12"	MD, M, MW	part sun to full shade	Beautiful forest groundcover with nearly round leaves. Strong ginger fragrance. Unusual flowers held on ground said to be pollinated by beetles. A must for forest restorations and shady perennial gardens.		3"	
Asclepias exaltata Poke Milkweed	White June to July	36-60"	MD, M	part sun to part shade	One of the most shade tolerant milkweeds it does best in dappled shade. It is also not as aggressive as common milkweed		3"	
Asclepias incarnata Swamp Milkweed	Lavender June to Aug	36-48"	M, MW, W	full sun to part shade	Showy plant of moist to wet habitats. Host plant for monarch butterflies, very good nectar source for butterflies and hummingbirds. Great for lake edges.	•	•	

Ascelpsias suberosa Unre to A General Contract Milkward Ascelpsias suberosa Unuter to Yee Sept 12-24" D, MD, M. Unit to Contract Milkward Ascelpsias suberosa Unuter to Sept Uniter to Weed Uniter to Milkward Whorled Milkward Aster Island Blue 24-30" MD, M, MW, Mill sun to part shade Shooth Aster Whorle Calcio Aster Mark noveweralgiare New England Aster Aster rowave-anglare New England Aster Aster rowave-anglare New England Aster Aster coverentargians New England Aster Aster ooverntargians New England Aster Aster poweranglare New Engla		Flowers	Ht.	Moisturo	Light	Description	2" Pot	4" Pot	6" Pot +
Common Milkweed  Orange  Drange  Drang	Asclepias svriaca					Description  Large leaved, robust plant known for being the larval host for	2	4	9
Subterly Weed   June to Sept						Monarch butterflies. Benefits many other pollinators as well.	•	•	
Asclepias verticilitate Whorled Milkweed July to Sept Sibre Whorled Milkweed July to Sept Sibre Whorled Milkweed July to Sept Sibre Short Asser in Selice July to Oct Aster felevis Smoch Asser July to Oct Aster felevis Sibre Aster menorphyllus Sibre Aster menorphyllus Sibre Aster menorphyllus Sibre Aster move-englide New England Aster Aug to Oct Aster oblongifolius Aster oblongifolius Aster oblongifolius Aster oblongifolius Aster oblongifolius Aster move-englide Aster felevis Aster oblongifolius Aster	Asclepias tuberosa		12-24"	D, MD, M	full to				
## the landscape but like other milkweed is terves as an important food source for Monarch cateprillars. Tolerant of drx soils.  ## Smooth Aster    July to Oct	Butterfly Weed				part sun		•	•	
sker laevis Smooth Aster  July to Oct Aster laevis Smooth Aster  July to Oct Aster laevis Smooth Aster  July to Oct Aster laevis Smooth Aster  White Age to Oct Aster  Aug to Oct Aster macrophyllus Blue Blue Blue Blue Blue Blue Blue Blue	· ·		12-24"	D, MD, M		l · · · · · · · · · · · · · · · · · · ·			
Aster leaves   Survey   20-30" MD, M, MW   full sun to Very showy vibrant blue-flowered peternals. Several cultivated   Smooth Aster   July to Oct   MD, M, MW   Full sun to Calico Aster   Aug to Oct   Sater macrophy/flus   Silve   Silve   Sater macrophy/flus   Silve   Sater mover-angliae   Silve   Aug to Oct   Aster mover-angliae   Silve   Sater mover-angliae   Silve mover-angliae   Silve   Sater mover-angliae   Silve mo	whoried Milkweed	July to Sept			sun	important food source for Monarch caterpillars. Tolerant of	•	•	
Aster Internations Calico Aster Aug to Oct Aster macrophyllus Bitue Bitue Bitue Aster macrophyllus Bitue Aster movae-angliae Aug to Oct Aster novae-angliae Aug to Oct Aster novae-angliae Aug to Oct Aster novae-angliae Aug to Oct Aster obligation Aster programicoles Aug to Oct Aster obligation Aster obligation Aster obligation Aster obligation Aster obligation Aster obligation Aster programicoles Upland White Aster Aug to Oct Aster programicoles Aster programicoles Upland White Aster Aug to Oct Aster programicoles Aster programicoles Upland White Aster Aug to Oct Aster programicoles As	Aster laevis		20-30"	MD, M, MW	full sun to	Very showy vibrant blue-flowered perennial. Several cultivated			
Aster macrophyllus Blue Blue Blue Blue Blue Blue Aster novae-angliee Aug to Oct W Aster novae-angliee Aug to Oct New England Aster Aug to Oct New England Aster Aug to Oct Aster novae-angliee Aug to Oct Aster novae-angliee Aug to Oct New England Aster Aug to Oct Aster oblingfiolius Aster punicus Sky Blue Aster Aug to Oct Aster unbolidius Upland White Aster Aug to Oct Aster unbolidius Plant oblingfiolius Aster punicus As						cutflower.	•	•	
Aster macrophyllus Big Leaf Aster  Aug to Oct  Aster novae-engliae New England Aster  Aug to Oct  Aster oblorgifolius  Aster oblorgifolius  Aster oblorgifolius  Aster parmicoides  Apart parmicoides  Upland White Aster  Aug to Oct  Aster parmicoides  White  Upland White Aster  Aug to Oct  Aster parmicoides  Upland White Aster  Aug to Oct  Aster parmicoides  White  Aug to Oct  Aster parmicoides  Upland White Aster  Aug to Oct  Aster unbellatus  Filal-topped Aster  Aug to Oct  Aster unbellatus  Baptisia alba  White  30-42° D, MD, M, MII Ill sun to  full shabe  White  Blue  30-42° M, MWW  part sun to  full shabe  Upland White Aster  Aug to Oct  Aster or particoides  Upland White Aster  Aug to Oct  Aster or particoides  Upland White Aster  Aug to Oct  Aster or particoides  Upland White Aster  Aug to Oct  Aster or particoides  Upland White Aster  Aug to Oct  Aster or particoides  Upland White Aster  Aug to Oct  Aster or particoides  Upland White Aster  Aug to Oct  Aster or particoides  Upland White Aster  Aug to Oct  Aster or particoides  Upland Touricoides  Upland T			24"	MD, M, MW					
Big Leaf Aster Algro Oct MW shade Broad heart-shaped leaves make an great, fast-growing	Calico Aster	Aug to Oct			snade		•	•	
Aster novae-angliae New England Aster New England Aster New England Aster Aug to Oct Aster obloragifolius Aster ob			8-18"	D, MD, M,	full sun to	· ·			
Aster oborgide New England Aster  Aug to Oct  Aster oborgide (New England Aster)  Aster oborgide (New	Big Leaf Aster	Aug to Oct		MW	shade		•	•	
New England Aster  Aug to Oct  Aster oblongifolius  Aster punicales  White  Aug to Oct  Aster punicales  White  Aug to Oct  Aster punicales  White  Aug to Oct  Aster punicales  Aug to Oct  Aster punicales  Aug to Oct  Aster punicales  White  Aug to Oct  Aster punicales  Aug to Oct  Aster punicales  White  Aug to Oct  Aster punicales  Aug to Oct  Aster punicales  Aug to Oct  Aster punicales  White  Aug to Oct  Aster punicales  Aug to Oct  Aster punicales  White  Aug to Oct  Aster punicales  White  Aug to Oct  Aster punicales  Aug to Carlongia  Bull to Oct  Aster punicales  Aug to Carlongia  Bul	Aster novae-analiae	Blue	24-48"	MD, M. MW.	full sun to				$\vdash$
Aster oblorating lates Aug to Oct Aster oblorations and any to Aug to Oct Aster punicates Aug to Oct	New England Aster					garden perennial and for wet lake-edge plantings. Nectar	•	•	
Accomance Aster Aug to Oct Sky Blue Aster Aug to Oct Sky Blue Aster Aug to Oct Aster optimizer restorations.  Aster optimizeriosides Sky Blue Aster Aug to Oct Aug D, MD, M I full to part sum Sky Blue Aster Aug to Oct Aster plannicoides Upland White Aster Plannicoides Upland White Aster Aug to Oct Aster Aug to Oct Aster Aug to Oct Aster Divide Aster Aug to Oct Aster Aug to Oct Aster Divide Aster Aug to Oct Aster Aug	Aster oblonaifolius	Violet	12-24"	D. MD	full to				
Aster colonitangiense Aug to Oct Aster parmicoides Upland White Aster Aug to Oct Aster parmicoides Aster Aug to Oct Aster parmicoides Upland White White White White Mode Upland White Aster Aug to Oct Aster parmicoides Upland White White White Mode Upland Vision to Aug White White White Mode Upland Vision to Aug Baptisia australis Blue June to July Dune to Sept Du				_,		throughout SE to NW MN and SW to NW WI. A must for	•	•	
Sky Blue Aster   Aug to Oct   Sun   among native wildlowers. Attracts loads of polinators.   Aster puniceus   Aug to Oct	Aster oolentangiense	Blue	24-36"	D, MD, M	full to part	<u> </u>			
Upland White Aster   Aug to Oct      Purple	Sky Blue Aster					-	•	•	
Excellent in short prairies.   Excellent in short prairies.   Aster puniceus   Purple   Aster puniceus   Aug to Oct   Aster umbellatus   Swamp Aster   Aug to Oct   Aster umbellatus   White   30-42"   M, MW, W   full sun to faster umbellatus   White   July to Aug   White   July to Aug   White   July to Aug   White   June to Aug   MW   Mile   June to Aug   MW   Mile   June to Aug   MW   Mile   June to Aug   MW   Aug   June to Aug   June to July   Aug   July to Oct   Aug   July to Oct   Aug   June to Sept   June to Aug   June to Aug   June to Sept   June to Aug   June to Aug   June to Sept   June to Aug   June to Aug   June to Aug   June to Sept   June to Aug   June to Aug   June to Aug   June to Sept   June to Aug   June to Aug   June to Aug   June to Sept   June to Aug   June to Sept   June to Aug   June to Sept   June to Sept   June to Sept   June t			12-24"	D, MD					
Swamp Aster   Aug to Oct   Part shade   International Part   Aster umbellatus   White   Aster umbellatus   Aster umbellatus   White   Aster umbellatus   Aster umbellatus   Aster umbellatus   Aster umbellatus   Aster umbellatus   Aster umbellatus   White   Aster umbellatus   A	Opiana White Aster	J			part suri	•			
Aster umbellatus Flat-topped Aster July to Aug  White Baptisia alba White June to Aug Baptisia alba White June to Sept Large Indiang Aster umbellatus Flat-topped Aster July to Aug  White July to Aug  Mill to part sun part sun part sun Stately perennial with glossy foliage and beautiful sky blue part sun part sun part sun part sun Stately perennial with glossy foliage and beautiful sky blue part sun flower stolleten garden perennial. Native south of our range but touch sterse. Excellent garden perennial. Native south of our range but touch sterse. Excellent garden perennial. Native south of our range but touch sterse. Excellent garden perennial. Native south of our range but touch sterse. Excellent garden perennial. Native south of our range but touch sterse. The sterse in NW WI.  Cathra palustris  Wall to part sun to shade  A biennial so grows quickly and flowers the first or second year and then self-seeds. Tall spires of violet-blue flowers good for quick color in restorations. A plant blant blant blant so part shade lance Leaf Coreopsis June to Aug  Coreopsis palmata  Yellow June to Aug  A self-all to wet lake edges and moist to wet prairies and moist to wet prairies and moist to wet part sin and text buttefly plant and cuttlower. Ranges throughout Will in all to part shade long restorations. A plant shade long restorations and a variety of soil conditions except very dry.  Blue July to Oct  Coreopsis palmata  Yellow June to Aug  A see of MW, W  Jull sun to part shade  Flatil sun to part sha	Aster puniceus		36-56"	MW, W			•		
Flat-topped Aster  July to Aug  white  Baptisia alba White Wide Indigo White Wide Mide Mide Wide Indigo White Wide Indigo Baptisia australis Blue June to July June to July Baptisia australis Blue June to July June to July Baptisia australis Blue June to July June to July Baptisia australis Blue June to July June to June to July  Baptisia australis Blue June to July June to June to July  Baptisia australis Blue June to July June to June to July  Baptisia australis Blue June to July  Blue False Indigo  June to June to July  Baptisia australis Blue June to July  June to June to July  Baptisia australis Blue June to July  June to Sept  Blue June to Sept  Blue July to Oct  Blue July to Oct  Campanula americana Tall Bellflower  July to Oct  Blue June to Sept  White July to Oct  Coreopsis Ianceolata Lance Leaf Coreopsis June to Aug  White Jule to Sept  June to Oct  Dalea aundida White June to Sept  Dicentra cuculiaria White Jule to Sept  Mice July to Sept  June to Sept  June to Oct  Dalea purpurea Purple Prairie Clover  Dicentra cuculiaria White July white July white July white July to Ose  June to Sept  Mice June to Sept  Ju			30-42"	M. MW. W	•	-			
Baptisia alba   White   June to Aug   June to July   Stately perennial with glossy foliage and beautiful sky blue part sun to flower clusters. Excellent garden perennial. Native south of our rance but touch enough for restorations here in NW WI.   Harbinger of spring. Usually associated with peaty ground and some flowing water, but can be planted on any wet oround. Great for pond achoes.   A biennial so grows quickly and flowers the first or second year and then self-seeds. Tall spires of violet-blue flowers good for quick color in partially shaded restorations and a variety of soil conditions. Scapet very dry.   Bulbe bell-shaped flowers atop delicate, grassy foliage. Great plant for restorations. Common throughout Wi in all dry habitats. Shade-tolerant.   Spikes of showy turtle-shaped flowers atop glossy dark green foliage. Nice for gardens and lake edge restorations.   Spikes of showy turtle-shaped flowers atop glossy dark green foliage. Nice for gardens and lake edge restorations.   Spikes of showy turtle-shaped flowers atop glossy dark green foliage. Nice for gardens and lake edge restorations.   Spikes of showy turtle-shaped flowers atop glossy dark green foliage. Nice for gardens and lake edge restorations.   Spikes of showy turtle-shaped flowers atop glossy dark green foliage. Nice for gardens and lake edge restorations.   Spikes of showy turtle-shaped flowers atop glossy dark green foliage. Nice for gardens and lake edge restorations   Spikes of showy turtle-shaped flowers atop glossy dark green foliage. Nice for gardens and lake edge restorations   Spikes of showy turtle-shaped flowers atop glossy dark green foliage. Nice for gardens and lake edge restorations   Spikes of showy turtle-shaped flowers atop glossy dark green foliage. Nice for gardens and lake edges restorations   Spikes of showy	Flat-topped Aster	July to Aug			full shade	of moist to wet lake edges and moist to mesic forests. Great	•	•	
Baptisia australis Blue False Indigo June to July  12-18" MW, W Arsh Marigold May to June  12-18" MW, W Arsh Marigold Marsh Marigold  12-18" MW, W Arsh Marigold  12-18" MW, W Arsh Marigold May to June  12-18" MW, W Arsh Marigold  13"  14"  15"  15"  16"  16"  16"  16"  16"  16	Baptisia alba	White	30-42"	D, MD, M,	full to				
Blue False Indigo  Zeltha palustris  Yellow Marsh Marigold  May to June  Blue Campanula americana Tall Bellflower  July to Oct  Ablemanula rotundifolia Harebell  Ablemanula Diute obert  White Diute Careopsis Janceolata Lance Leaf Coreopsis June to Oct  Coreopsis Janceolata Lance Leaf Coreopsis June to Oct  Ablemanula  Yellow June to Oct  July to Oct  July to Oct  Ablemanula  Yellow June to Sept  June to Oct  July to Oct  July to Oct  Ablemanula	White Wild Indigo	June to Aug		MW	part sun		•	•	
Campanula americana Tall Bellflower  Blue July to Oct  MD, M, MW  Faringer of spring. Usually associated with peaty ground and some flowing water, but can be planted on any wet around. Great for pond edges.  A biennial so grows quickly and flowers the first or second year and then self-seeds. Tall spires of violet-blue flowers good for quick color in partially shaded restorations and a variety of soil conditions except very dry.  Campanula rotundifolia Harebell  White July to Oct  MD, M, MW  Part sun to part shade  year and then self-seeds. Tall spires of violet-blue flowers good for quick color in partially shaded restorations and a variety of soil conditions except very dry.  Blue bell-shaped flowers atop delicate, grassy foliage. Great plant for restorations. Common throughout WI in all dry habitats. Shade-tolerant.  Chelone glabra  Turtlehead  White July to Oct  Widespread in MN and WI.  Coreopsis lanceolata Lance Leaf Coreopsis  Coreopsis palmata  Prairie Coreopsis  Dalea candida  White June to Sept  16-30"  D, MD, M  full sun to part shade  Turtlehead  Dalea candida  White June to Sept  MW, W  Fullow June to Oct  D, MD, M  full to Dense 2" clusters of small white flowers atop grayish fermy foliage. Were arready and the self-seeds. Tall spires of violet-blue flowers good for quick color in partially shaded restorations and a variety of soil conditions except very dry.  Blue bell-shaped flowers atop glossy dark green foliage. Nice for gardens and lake edge restorations.  Widespread in MN and WI.  Wiry stems bearing yellow daisies. Easy to grow on toughest, driest soil types. Very good butterfly plant.  Coreopsis palmata  Prairie Coreopsis  Dalea candida  White June to Oct  D, MD, M  full to Dense 2" clusters of small white flowers atop grayish fermy foliage. Very attractive perennial for prairie and savanna restorations as well as perennial gardens.  Dalea candida  White June to Sept  D, MD, M  full to Similar to Dalea candida but with purple flowers. Both species are excellent butterfly plants. Used as tea	Baptisia australis		30-42"	M, MW					
Caltha palustris Marsh Marigold May to June May to Oct May and then self-seeds. Tall spires of violet-blue flivers second year and then self-seeds. Tall spires of violet-blue flivers second year and then self-seeds. Tall spires of violet-blue flivers May to June to Sept  May to June to Sept May to June May to Sept May to June Ma	Blue False Indigo	June to July			part sun	· ·	•	•	
Campanula americana Tall Bellflower  Blue July to Oct July to Oct July to Oct  Blue July to Oct July to Oct  Campanula rotundifolia Harebell Harebell Harebell  White July to Oct  Coreopsis lanceolata Lance Leaf Coreopsis June to Aug June to Oct  White June to Oct  Ablennial so grows quickly and flowers the first or second year and then self-seeds. Tall spires of violet-blue flowers good for quick color in partially shaded restorations and a variety of soil conditions except very dry.  Chelone glabra Turtlehead  White July to Oct  White July to Oct  Coreopsis lanceolata Lance Leaf Coreopsis  June to Aug  Torrice Coreopsis  June to Oct  Torrice Coreopsis  Torrice Coreopsis  June to Oct  Torrice Coreopsis  Torrice Coreopsis  June to Oct  Torrice Coreopsis  Torrice Coreops	'		12-18"	MW, W	l ·	Harbinger of spring. Usually associated with peaty ground			
Campanula americana   Tall Belliflower   July to Oct   Sept   July to Oct   Sept   July to Oct   J	Marsh Marigold	May to June			shade	, ,		3"	
Good for quick color in partially shaded restorations and a variety of soil conditions except very dry.	Campanula americana	Blue	36-60"	MD, M, MW	part sun to				
Variety of soil conditions except very dry.   Variety of soil conditions except very dry.	Tall Bellflower	July to Oct			part shade	The state of the s	•		
Campanula rotundifolia   Harebell   June to Sept   16-24"   D, MD   full sun to part shade   Plant for restorations. Common throughout WI in all dry habitats. Shade-tolerant.   Spikes of showy turtle-shaped flowers atop glossy dark green foliage. Nice for gardens and lake edge restorations.   Wirestendand or part shade   Wilestendand or part shade   Wilestendand or part shade   Wilestendand or part shade   Vellow June to Aug   D, MD   full sun to part shade   Vellow June to Oct   D, MD, M   full sun to part shade   Vellow June to Oct   D, MD, M   full sun to part shade   Vellow June to Oct   D, MD, M   full sun to part shade   Vellow daisies above ferny foliage. Native of driest prairies and bluff communities throughout southern 2/3 of MN and WI.   Oct   Good for butterflies.   Oct   Good for butterflies.   Oct   O									
habitats. Shade-tolerant.	Campanula rotundifolia	Blue	16-24"	D, MD	full sun to				
Chelone glabra Turtlehead  White July to Oct  Vellow June to Aug  Coreopsis palmata Prairie Coreopsis  Dalea candida White Prairie Clover White Prairie Clover  Dalea purpurea Purple Prairie Clover  Divide Coreopsis  White June to Sept  Similar to Dalea candida but with purple flowers. Both species are excellent butterfly plants. Used as tea by Native  Americans.  White June to Sept  Similar to Dalea candida but with purple flowers. Both species are excellent butterfly plants. Used as tea by Native  Americans.  Spring blooming woodland wildflower of mesic to moist forests  June to Sept  Spring blooming woodland wildflower of mesic to moist forests	Harebell	June to Sept			part shade		•	3"	
Turtlehead  July to Oct  part shade  foliage. Nice for gardens and lake edge restorations.  Widespread in MN and WI.  Wiry stems bearing yellow daisies. Easy to grow on toughest, driest soil types. Very good butterfly plant.  Coreopsis palmata Prairie Coreopsis  June to Oct  Part shade  Turtlehead  Yellow June to Aug  16-30"  D, MD, M  full sun to part shade  Prairie Coreopsis  Dalea candida White Prairie Clover  Wiry stems bearing yellow daisies. Easy to grow on toughest, driest soil types. Very good butterfly plant.  Yellow June to Oct  Purple June to Sept  Purple June to Sept  D, MD, M  full to part sun  Furple Prairie Clover  White  Furple Prairie Clover  Furple Prairi	Chelone alabra	White	24-36"	MW. W	full sun to				
Coreopsis lanceolata Lance Leaf Coreopsis  June to Aug  Yellow June to Oct  Yellow June to Oct  Yellow June to Oct  Yellow June to Oct  In June to Oct  Yellow June to Oct  Yellow June to Oct  June to Oct  In June to Oct  Yellow daisies above ferny foliage. Native of driest prairies and bluff communities throughout southern 2/3 of MN and WI. Good for butterflies.  Dense 2" clusters of small white flowers atop grayish ferny foliage. Very attractive perennial for prairie and savanna restorations as well as perennial gardens.  Dicentra cucullaria  Yellow June to Oct  Yellow daisies above ferny foliage. Native of driest prairies and bluff communities throughout southern 2/3 of MN and WI.  Yellow daisies above ferny foliage. Very attractive perennial for prairie and savanna restorations as well as perennial gardens.  Similar to Dalea candida but with purple flowers. Both species are excellent butterfly plants. Used as tea by Native  Americans.  Note of the top of the story of driest prairies and bluff communities throughout southern 2/3 of MN and WI.  Yellow daisies above ferny foliage. Native of driest prairies and bluff communities throughout southern 2/3 of MN and WI.  Yellow daisies selve grow on toughest, driest soil types. Very good butterfly plants.  Yellow daisies selve grow on toughest, driest soil types. Very good butterfly plants.  Yellow daisies selve grow on toughest, driest soil types. Very good butterfly plants.	_			,		foliage. Nice for gardens and lake edge restorations.	•		
Lance Leaf Coreopsis  June to Aug  Coreopsis palmata Prairie Coreopsis  Dalea candida White Prairie Clover  Dalea purpurea Purple Prairie Clover  Dicentra cucullaria  White  June to Aug  Part shade  Dyert shade  Dyert shade  Dyert shade  Dyellow daisies above ferny foliage. Native of driest prairies and bluff communities throughout southern 2/3 of MN and WI.  Good for butterflies.  Dense 2" clusters of small white flowers atop grayish ferny foliage. Very attractive perennial for prairie and savanna restorations as well as perennial gardens.  Similar to Dalea candida but with purple flowers. Both species are excellent butterfly plants. Used as tea by Native  Americans.  Dicentra cucullaria  White  16-30"  D, MD, M  full to part sun  Furple June to Sept  M, MW  Full to part Spring blooming woodland wildflower of mesic to moist forests  3"	Coroonsia langualata	Vollow	24.26"	D MD	full oun to				<u> </u>
Coreopsis palmata Prairie Coreopsis  Une to Oct  Prairie Coreopsis  Prairie Coreopsis  Prairie Coreopsis  Prairie Coreopsis  Une to Oct  D, MD, M  Full sun to part shade  Good for butterflies.  Dense 2" clusters of small white flowers atop grayish ferny foliage. Very attractive perennial for prairie and savanna restorations as well as perennial gardens.  Purple June to Sept  D, MD, M  Full to part sun  Fu	· · · · · · · · · · · · · · · · · · ·		24-30	טואו ,ט			•	•	
Good for butterflies.  Dalea candida White Prairie Clover Unue to Sept White Prairie Clover Unue to Sept Unue	Coreopsis palmata	Yellow	16-30"	D, MD, M	full sun to	Yellow daisies above ferny foliage. Native of driest prairies			
Dalea candida White June to Sept White Prairie Clover White Prairie Clover  Dense 2" clusters of small white flowers atop grayish ferny foliage. Very attractive perennial for prairie and savanna restorations as well as perennial gardens.  Purple Purple June to Sept Dense 2" clusters of small white flowers atop grayish ferny foliage. Very attractive perennial for prairie and savanna restorations as well as perennial gardens.  Similar to Dalea candida but with purple flowers. Both species are excellent butterfly plants. Used as tea by Native Americans.  Dicentra cucullaria White 12" M, MW full to part Spring blooming woodland wildflower of mesic to moist forests 3"	Prairie Coreopsis	June to Oct			part shade	· ·	•	•	
White Prairie Clover  June to Sept  part sun  foliage. Very attractive perennial for prairie and savanna restorations as well as perennial gardens.  Purple June to Sept  Purple June to Sept  16-30"  D, MD, M  full to part sun  foliage. Very attractive perennial for prairie and savanna restorations as well as perennial gardens.  Similar to Dalea candida but with purple flowers. Both species are excellent butterfly plants. Used as tea by Native Americans.  Dicentra cucullaria  White  12"  M, MW  full to part Spring blooming woodland wildflower of mesic to moist forests  3"	Dalea candida	White	16-30"	D, MD, M	full to				
Dalea purpurea       Purple       16-30"       D, MD, M       full to part sun       Similar to Dalea candida but with purple flowers. Both species are excellent butterfly plants. Used as tea by Native Americans.         Dicentra cucullaria       White       12"       M, MW       full to part       Spring blooming woodland wildflower of mesic to moist forests       3"	White Prairie Clover	June to Sept			part sun	foliage. Very attractive perennial for prairie and savanna	•	•	
Purple Prairie Clover  June to Sept  part sun  are excellent butterfly plants. Used as tea by Native  Americans.  Dicentra cucullaria  White  12"  M, MW  full to part  Spring blooming woodland wildflower of mesic to moist forests  3"	Dalea purpurea	Purple	16-30"	D. MD. M	full to				<del>                                     </del>
Dicentra cucullaria White 12" M, MW full to part Spring blooming woodland wildflower of mesic to moist forests 3"	Purple Prairie Clover						•	•	
	Diagratus augustus vie	Mhito	40"	NA NAVA/	full to man				
			12"	IVI, IVIVV				3"	

	Flowers	Ht.	Moisture	Light	Description	2" Pot	4" Pot	6" Pot +
Dadaaanthaan maadia	White	18"	MD, M, MW			2	4	9
Dodecantheon meadia Shooting Star	May to June	10	INID, INI, INIV	sun	Cute plant with basal leaves and a leafless flowerstalk with shooting star-like white to pink flowers. Dry prairies, savannas, and woodlands on droughty soils. Rare in extreme SE MN and scattered in SW WI		3"	
Echinacea angustifolia Narrow-Leaf Purple Coneflower	Lavender July to Aug	24-36"	D, MD	full to part sun	Said to be the best of the purple coneflowers for medicinal purposes. Native in Western MN and further west, but hardy in this area. Very attractive with narrow leaves.	•	•	
Echinacea pallida Pale Purple Coneflower	Lavender July to Aug	24-36"	D, MD, M	full to part sun	Only echinacea species definitely native to WI. Nice narrow foliage. Heads have reflexed rays ("petals"). Can't beat these for butterflies.	•		
Echinacea purpurea Purple Coneflower	Purple July to Aug	24-36"	MD, M, MW	full to part sun	Spectacular display of broad-headed purple daisies. Great for restorations and perennial gardens.	•	•	
Eryngium yuccifolium Rattlesnake Master	White July to Oct	28-42"	MD, M, MW	full to part sun	Exotic-looking plant that resembles a desert yucca. Bristly white heads and whitish foliage make it a great background perennial as well as a tough fast grower for restorations.	•	•	
Eupatorium maculatum Spotted Joe-Pye Weed	Purple July to Sept		MW, W	full sun to part shade	Good lake edge perennial with broad flat-topped clusters of purple heads. Excellent butterfly plant. Ranges throughout MN and WI.	•	•	
Eupatorium perfoliatum Boneset	White June to Aug		MW, W	full sun to part shade	Nice lake edge plants with flat-topped cluster of white heads. Very nice purplish foliage with opposite fused leaves. Used medicinally to hasten healing of bones. Ranges throughout MN and WI.	•		
Eupatorium purpureum Purple Joe Pye Weed	Purple July to Sept	24-36"	MD, M, MW	part sun to shade	Moist to moderately dry savannas and forests through southern 2/3 of WI and eastern 1/2 of MN. Attractive broad clusters of purple heads. Excellent for attracting butterflies.	•	•	
Euthamia graminfolia Grass-leaved Goldenrod	Yellow July to Sept	48"	MD, M, MW, W	Full sun	This plant is not really a goldenrod but is an excellent source of pollen for bees. Can be somewhat spreading and aggressive but is very adaptable to a variety of soil moisture conditions and good for restorations.	•	•	
Fragaria vesca Woodland Strawberry	White May to June	6-10"	MD, M, MW	full sun to shade	Attractive groundcover that grows rapidly. A must for the woodland restoration. Sweet tasting red fruits in June are a bonus. Ranges throughout the world in northern latitudes including all of MN and WI.		3"	
Fragaria virginiana Wild Strawberry	White May to June	6"	D, MD, M, MW	full sun to shade	Stoloniferous spreading groundcover bears delicious fruit. Found throughout MN and WI ranging from prairies to forests. Widely adapted to dry, moist or even wet soils.	•	•	
Galium boreale Northern Bedstraw	White June to July	16-24"	MD, M, MW, W	full sun to full shade	Common understory species of northern forests in MN & WI. Easy to grow on mesic to dry soils.	•	•	
Geranium maculatum Wild Geranium	Pink May to June	18"	MD, M	full to part shade	Tolerant to all soils and woodland conditions except deeply shaded situations. Nice perennial geranium with attractive foliage that turns red in fall.	•	3"	
Geum triflorum Prairie Smoke	Red May to June	6-12"	D, MD, M, MW	full to part sun	Awesome early-flowering prairie wildflower, great for sandy site restorations and for rock gardening. Hairy fruits remain attractive throughout the year.	•		
Helenium autumnale Sneezeweed	Yellow July to Oct	30-48"	MW, W	full sun to shade	Sneezeweed won't make you sneeze - it was used as an expectorant by Native Americans. Gaillardia -like heads in abundance. Good butterfly plant.	•		
Helianthus strumosus Woodland Sunflower	Yellow July to Oct	36-48"	MD, M, MW	full sun to part shade	Very common woodland sunflower seen at the edge of forests and ditches. Variable but showy with terminal yellow heads. Birds eat seeds. Excellent soil stabilizer.		3"	
Heliopsis helianthoides False/Early Sunflower	Yellow June to Oct	24-42"	MD, M, MW	full to part sun	Widespread on dry to wet soils. Long flowering period. Suited to all but very shady restoration sites. Commonly used as a garden perennial.	•	•	
Hepatica acutiloba Sharp-lobed Hepatica	White, Pink, Blue April to May	3-6"	MD, M	part sun to part shade	One of the earliest blooming wildflowers you will find. A small tuft of basal leaves are mottled green to reddish brown and persist through the winter.		3"	
Heuchera richardsonii Alum Root	White May to July		D, MD, M, MW	full sun to part shade	Common through most of WI and MN. Attractive, tough ground cover well suited to dry soils. Quite shade-tolerant. Usually associated with rock outcrops in prairies, woodlands, and forests.	•	•	
<i>Iris versicolor</i> Northern Blue Flag Iris	Blue June to July		M, MW, W	full to part sun	Common on lake margins throughout northern WI and MN. Very attractive, forming large clumps. Some consider this to be synonymous with <i>Iris virginica</i> .	•	•	
Liatris aspera Rough Blazing Star	Purple July to Sept	18-36"	D, MD, M	full to part sun	Beautiful purple spikes. Most common liatris of dry, sandy prairies throughout MN and WI. Excellent butterfly plant.	•	•	

	Flowers	Ш	Moiotura	Light	Description	2" Pot	4" Pot	6" Pot +
Liatris ligulistylis	Purple	Ht.	Moisture MD, M, MW		Description   Rosy heads appear in abundance on tall spikes. Beautiful	7	4	9
Northern Plains Blazing Star	July to Aug			part sun	plant among the best to attract butterflies. OK on wet and dry soils.		•	
Liatris pychnostachya Thick Spike Blazing Star	Purple July to Sept	24-36"	M, MW, W	full to part sun	Dense purple spikes. Good for moist and wet restoration sites. Very showy, good butterfly plant. Ranges throughout prairie regions of MN and WI.	•	•	•
Liatris spicata Marsh Blazing Star	Purple July to Aug	24-36"	M, MW, W	full to part sun	Most common liatris for perennial gardening. Grows well in any condition. Perfect pollinator plant. Native into SE WI. Moist to wet soils.		•	
Lilium michiganense Michigan Lily	Orange June to Aug	36-60"	M, MW, W	full to part sun	Orange flowers with black spots make this rare native lily very similar to the classic garden Turk's Cap Lily.		3"	
Lobelia cardinalis Cardinal Flower	Red July to Oct	36-48"	MW, W	full sun to part shade	Showy spikes of red flowers are the best for attracting hummingbirds. Suited to moist soils particularly along waterways. Native through most of WI.		•	
Lobelia siphilitica Great Blue Lobelia	Blue July to Oct	20-30"	M, MW, W	full sun to part shade	Widely adapted to wet and moist soils. Vivid blue and white- striped flowers in terminal spikes. Compact and well suited to perennial gardens.		•	
<i>Mertensia virginiana</i> Virginia Bluebells	Blue May to June	18-24"	M, MW	part to full shade	Clusters of bell-shaped blue flowers in spring. Nice woodland wildflower. Dies back by midsummer, but are plantable while dormant.		3"	
<i>Mimulus ringens</i> Monkeyflower	Lavender July to Sept		MW, W	full sun to part shade	Lavender spikes of 1" snapdragon flowers that look like small monkeys. Compact plants. Excellent for lakescaping.	•	•	
<i>Monarda fistulosa</i> Bergamot	Lavender July to Aug	36-48"	D, MD, M, MW	full sun to part shade	One of the most common and widespread native wildflowers. Good in herbal teas. Used medicinally by Native Americans. Moist to dry soils.		•	
Monarda punctata Dotted Mint	Lavender June to Sept	18-30"	D, MD	full to part sun	Fragrant herb with showy dotted multicolored leaves below flowers. Good on dry soils. Used medicinally and in herbal teas. Nice cutflower.		•	
Penstemon digitalis Smooth Penstemon	White June to July	24-30"	MD, M	full sun to part shade	1.5" white tubular flowers in tall spikes. Good garden perennial and cutflower. Tolerant of dry and mesic soils and a variety of light conditions.		•	
Phlox divaricata Woodland Phlox	Blue June to July	12-18"	MD, M, MW	part sun to shade	Beautiful fragrant spring wildflower occurring in forests ranging from mesic to floodplain. Excellent for shady restorations and gardens.		•	
Phlox pilosa Prairie Phlox	Pink May to June	18-24"	D, MD, M, MW	full to part sun	Broad heads of deep pink flowers. Very nice wildflower for restorations and perennial gardens.		•	
Physostegia virginiana Obedient Plant	Pink July to Oct	24-36"	M, MW, W	full sun to part shade	Popular garden perennial native to moist prairies, lakeside and streamside wetlands.	•		
Polemonium reptans Jacob's Ladder	Blue May to June	18"	MD, M, MW	part sun to full shade	Beautiful spring woodland wildflower. Ferny leaves resemble a ladder. A must for woodland gardens and restorations.	•	•	
Polygonatum biflorum Solomon's Seal	White May to June		MD, M	part sun to full shade	A very attractive plant with graceful arching stems that add character and charm to any shaded garden or restoration.			٠
Ratibida pinnata Yellow Coneflower	Yellow July to Oct	24-52"	MD, M	full to part sun	One of the more common native prairie wildflowers. Very attractive to butterflies. Greyish foliage makes this a nice contrast plant.	•	•	
Rudbeckia fulgida Orange Coneflower	Orange- Yellow July to Oct	24-36"	MD, M, MW	full to part sun	Beautiful compact coneflower with deep green foliage and 3" orange-yellow heads with brown centers. Parent of 'goldsturm', one of the most popular garden perennials.		•	
Rudbeckia hirta Black Eyed Susan	Yellow June to Oct	24-36"	MD, M, MW	full to part sun	Very common short-lived perennial occurring in a wide range of habitats. Produces lots of color the first year.		•	
Rudbeckia laciniata Wild Goldenglow	Yellow July to Sept	36-48"	M, MW	full sun to part shade	Found throughout WI, primarily in swamp habitats and along lakes and streams, but also in prairies. Good for moist, shaded restoration sites.			
Rudbeckia subtomentosa Sweet Black Eyed Susan	Yellow July to Sept	24-36"	MD, M, MW	full to part sun	Very attractive yellow heads in profusion atop deep green glossy leaves. Good for restorations and perennial gardens. Native into SW WI.	•	•	
Rudbeckia triloba Brown Eyed Susan	Yellow July to Oct	24-36"	MD, M, MW	full sun to part shade	Outstanding coneflower with numerous heads which are smaller than other species, but more abundant. Very nice foliage. Should be grown more widely in perennial beds. Mesic to wet soils.	•	•	
Sanguinaria canadensis Bloodroot	White April to June	8"	MD, M, MW	full to part shade	Spring blooming woodland wildflower found throughout the forest supporting regions of MN and WI. Single, jagged umbrella shaped leaf is accompanied by a single, large, showy white flower.		3"	

	Flowers	Ht.	Moisture	Light	Description	2" Pot	4" Pot	6" Pot +
Silene regia	Red	18-24"	MD, M	full sun to	Deep red tubular flowers flaring out into a star. Excellent			
Royal Catchfly	June to Oct			part shade	hummingbird plant and garden perennial. Native just south of	•	•	
					MN and WI, but hardy in the north.			
Silphium perfoliatum	Yellow	48-72"	M, MW	full sun to	Impressive sunflower with numerous broad yellow heads.			
Cup Plant	July to Sept			part shade	Unusual opposite leaves fuse to make cups which collect	•		
	,				water for bird bathing. Birds love it!			
Smilacina racemosa	White	18-24"	MD, M, MW	full to part	Two-foot tall leafy stemmed lily with a terminal cluster of small			
False Solomon Seal	May to June		, ,	shade	white flowers, each of which becomes a red berry. Found		3"	
	,				throughout forest supporting regions of MN and WI.			
Solidago flexicaulis	Yellow	24"	MD, M, MW	part to	Good woodland perennial for restorations. Unique zig-zag			
Zig Zag Goldenrod	Aug to Sept			full shade	stems with scattered yellow heads. Very different than most			
					goldenrods. Moist to mesic soils.			
Solidago nemoralis	Yellow	24-36"	D. MD	Sun to part	One of the most attractive goldenrods, it has vibrant golden-			
Gray Goldenrod	Aug to Oct			shade	yellow flowers that are a great source of pollen for bees and			
Gray Golderii Ga	riag to out			Silade	butterflies right up to frost. It is also one of the most shade-	•		
					tolerant goldenrods.			
Solidago rigida	Yellow	24-36"	D, MD, M,	full to	Another goldenrod worthy of growing for its very attractive			
Stiff Goldenrod	Aug to Sept		MW	part sun	foliage. Topped with a broad, flat cluster of yellow heads.		١.	
Cim Coldonioa	rag to copt			partouri	Adapted to mesic and moist soils. Good garden perennial.			
Solidago speciosa	Yellow	24-36"	D, MD, M	full sun to	Highly recommended for restorations and for perennial			
Showy Goldenrod		24-30	D, IVID, IVI	part shade	gardens. Very showy tall heads above glossy green foliage.		١.	
Showy Goldeniod	Aug to Sept			part snaue		•	•	
T	3.841.24	0.4"	145 14 144		Great butterfly plant.			
Thalictrum dioicum	White	24"	MD, M, MW	part to	Nice woodland wildflower for all but the driest sandy soils.		۵,	
Early Meadow Rue	May to June			full shade	Pretty flower heads held above columbine-like foliage. Nice in		3"	
Tradagaentia obiensia	Blue	04.40"		full sun to	the shaded perennial garden.			
Tradescantia ohiensis		24-48"	D, MD, M,		Forms large clumps creating a very nice display of 1" flowers			
Ohio Spiderwort	June to Oct		MW	part shade	in heads on top of each stem. Great for restorations and	•	١.	
	3.40.24				gardens. Moist to dry soils.			
Trillium grandiflorum	White	8-12"	M, MW	part shade	Spectacular 2" white 3-petaled flowers in May. This harbinger			
Trillium	May			to shade	of spring will let you know that winter is finally over!		3"	
					Widespread in WI and MN forests.			
Verbena hastata	Blue	36-48"	M, MW, W	full sun to	Tall spikes of blue flowers really attract butterflies. Great for			
Blue Vervain	June to Sept			part shade	shoreland restorations on mesic to saturated soils.	•		
					Throughout MN and WI.			
Verbena stricta	Blue	24-36"	D, MD	full sun to	Spikes of blue flowers above greyish foliage. Adapted to dry,		١.	
Hoary Vervain	June to Sept			part shade	sandy soils. Good butterfly plant.	•	Ι΄.	
Vernonia fasciculata	Rose-Pink	24-42"	M, MW	full sun to	Broad, flat clusters of pink heads to 5" across. Very attractive			
Ironweed	July to Sept			part shade	to butterflies. Good in moist to wet soils for lake edge	•		
					landscaping. Wide ranging in MN and WI.			
Veronicastrum	White	36-54"	MD, M, MW	full sun to	Several tall showy spikes on top of each stem make this an	1		
virginianum	June to Sept			part shade	excellent tall garden perennial. Attracts butterflies. Good in		١.	
Culver's Root	Same 10 Copt				most soils except externely dry.			
Viola pedatifida	Blue	4"	MD, M	full sun to	1.5" blue flowers above deeply clawed foliage. Very nice for		1	
Prairie Violet		7	ואוט, ואו		edging restoration sites on dry to mesic soils. Also nice for		3"	
i idilie violet	May to June			part shade			٦	
7:-:	Vallan	40.00"	MD M MAY	full accepta	rock gardening. Common in prairie regions of MN and WI.		<u> </u>	
Zizia aurea	Yellow	18-30"	MD, M, MW	full sun to	Early-flowering plant of mesic to moist habitats with 3" flat-			
Golden Alexanders	May to July			part shade	topped heads of yellow flowers. Seed heads that follow are	•	Ι.	
ĺ					attractive throughout the year.	l		

	Flowers	Ht.	Moisture	l iaht	Description	2" Pot	4" Pot	6" Pot +
Grasses, se				_	Description	N	4	9
Andropogon gerardii Big Bluestem	Purplish Green July to Sept	36- 72+"	D, MD, M, MW	full sun to part shade	One of the most dominant tall prairie grasses. Purplish foliage creates a nice effect along with turkey-foot-like flower head. Dry to moist habitats.	•		
Bouteloua curtipendula Side Oats Grama	Red-Green July to Sept	18-30"	D, MD, M	full sun to part shade	Very common species of dry to mesic habitats. Stamens are red, making them beautiful cutflowers when fresh. A must for any open dry site restoration.	•		
Bouteloua gracilis Blue Grama	Purplish Green July to Sept	12"	D, MD	full to part sun	Native through the central Great Plains into western WI. Very good lawn alternative. Low growing, forming attractive clumps.	•		
Calamagrostis canadensis Blue Joint Reedgrass	Straw July to Aug	36-48"	M, MW, W	full sun to part shade	Fine-leaved grass forms a dense cover on lake margins. One of the grasses displaced by reed canary grass and a good one to plant after nuking RCG. Purplish foliage.			
Carex bebbii Bebb's Sedge	Green May to June	24-36"	M, MW	full sun to part shade	Forms dense clumps on moist to saturated soil. Very common in most wetland areas throughout MN and WI.	•		
Carex bicknellii Bicknell's Sedge	Tan May to June		MD, M, MW	full to part sun	Sedge adapted to habitats ranging from dry to moist, usually on rocky, gravelly soil. Forms nice clumps. Found throughout southern 2/3 of MN and WI.	•		
Carex comosa Bottlebrush Sedge	Green May to June		MW, W	full sun to part shade	Beautiful clumps of yellowish stems on lake margins. Nodding flower spikes look like a bottlebrush. Excellent for restorations and water gardens.	•		
Carex crinita Caterpillar Sedge	Green May to June		MW, W	full sun to part shade	Clump-forming. 4" nodding spikes resemble a caterpillar. Very nice for wet site restorations and for water gardens. Excellent for control of shoreland erosion.	•		
Carex gracillima Graceful Sedge	Green May to June	18-30"	MW, W	full sun to full shade	Clump forming plant of mesic to moist forests throughout MN and WI. Attractive nodding flower spikes.	•	•	
Carex hystricina Porcupine Sedge	Green May to June	24-42"	MW, W	full sun to part shade	Very similar to <i>Carex comosa</i> . Very attractive plants for shoreland restorations and water gardening.	•		
Carex lupulina Hop Sedge	Green May to June	24-42"	MW, W	full sun to part shade	Flower spikes resemble hops clusters. Stout-stemmed clump former of a wide variety of wet habitats, including shady swamps. Very good for shady rain gardens.	•		
Carex pennsylvanica Pennsylvania Sedge	Brown May	6-8"	D, MD, M	full sun to full shade	Outstanding, low, clump-forming graminoid with yellow-green foliage. Forms extensive stands in northwoods forests that are extremely attractive. A must for forest restorations in northern MN and WI.	•		
Carex stipata Awl Fruit Sedge	Green May to June	16-40"	MW, W	full sun to full shade	Common, coarse clump forming sedge found in a wide variety of wet habitats throughout MN and WI. Very good fast grower. Shade tolerant species.	•		
Carex stricta Tussock Sedge	Green May to June	24-36"	MW, W	full sun to part shade	Forms dense clumps of closely clustered fine stems in a variety of wetlands and lake margins. Good shoreland and water garden plant.			
Carex vulpinoidea Fox Sedge	Green May to June	18-30"	MD, M, MW, W	full sun to part shade	Beautiful clump-forming sedge of lake and wetland margins. Very attractive multi-tiered flower spikes persist through the summer and into the fall.	•		
Elymus hystrix (patula) Bottlebrush Grass	Green June to July	36-48"	MD, M	full sun to full shade	Clump forming grass with tall spikes topped with an infloresence that looks like a bottlebrush. A must for forest restorations and shady perennial gardens.	•		
Glyceria canadensis Rattlesnake Manna Grass	Green July to Aug	36"	MW, W	full to part shade	A cool-season grass with distinct showy seed heads. Great for addint texture and interest to wet restorations.	•		
Glyceria striata Fowl Manna Grass	Green July to Aug	24-36"	M, MW, W	full sun to full shade	Low clump forming grass common next to small pools in swamps, in moist to seasonally wet forests, and along waterways.	•		
Hierochloe odorata Sweet Grass	Green May to June		M, MW, W	full sun to part shade	Widespread in MN and WI mostly on wetland margins on moist to wet soils. Sacred plant for Native Americans, used for braiding and basket-weaving. Revered for its sweet, aromatic vanilla-like fragrance.	•		
Juncus effusus Soft Rush	Brown July to Aug	24-48"	M, MW, W	full sun to part shade	Deep green wirey stems in large clumps along waterways and in a variety of wetlands. Evergreen plants remain attractive into winter.			
Juncus tenuis Path Rush	Brown June to July	to 16"	D, MD, M, MW	full sun to part shade	Very common, widespread plant of wet places throughout MN and WI. Nice for restoration site margins. Easy to grow. Forms attractive clumps.			

	Flowers	Ht.	Moisture	Light	Description	2" Pot	4" Pot	6" Pot +
Koeleria macrantha June Grass	Green June to July	12-24"		full sun to part shade	Clump-forming grass, most commonly associated with dry sandy soils. Low-grower suitable for edging native restoration plantings.	•	•	
Panicum virgatum Switch Grass	Purple June to Oct	36-96"	D, MD, M, MW	full to part sun	Fast growing clump former with attractive open infloresence. Suited to soils ranging from mesic to wet. Good for steep banks and slopes.	•		
Schizachyrium scoparium Little Bluestem	Amber July to Sept		D, MD, M	full sun to part shade	Fabulous ornamental grass native to dry, sandy to mesic habitats throughout the region. Amber foliage maintains color through winter. Forms neat clumps.	•		
Scirpus atrovirens Green Bulrush	Brown June to July	36-48"	MW, W	full sun to part shade	Coarse clump forming plant with nice clusters of small brown heads. Suitable for wet shoreland zones.	•		
Scirpus cyperinus Wool Grass	Brown June to July		MW, W	full sun to part shade	Coarse clump former with brown fuzzy heads that appear wooly. One of the most common lake edge species throughout the region.	•		
Sorghastrum nutans Indian Grass	Amber July to Sept		D, MD, M	full sun to part shade	Clump-forming grass of dry, sandy to moist habitats. Very ornamental and suitable for the back of perennial borders. Great plant for a wide range of habitats.	•		
Spartina pectinata Prairie Cordgrass	Yellow Aug to Oct	48-72"	M, MW, W	full to part sun	Rhizomatous grass forming large clumps in sandy soil, usually near the water table, but sometimes under dry conditions.  Common shoreland plant in northern WI and MN.	•		
Sporobolus heterolepis Prairie Dropseed	Green Aug to Oct	18-36"	D, MD, M, MW	full to part sun	Clump former with open, fine sprays of flowers. Popular as a garden perennial. Native to dry and mesic prairies throughout the region.	•		
Ferns Adiantum pedatum Maidenhair Fern		24"	MD, M, MW	part to full shade	Graceful stems with soft highly divided leaves in a U-shaped pattern. Slow-starting but eventually forms nice clumps. Moist to wet forests.		3"	
Athyrium felix-femina Lady Fern		36"	M, MW	part to full shade	Semi-spreading lime green highly divided leaves make this a very useful ground cover for shady restorations and gardens.		3"	
Dryopteris marginalis Eastern Wood Fern		20"	MD, M, MW	part to full shade	Blue-green highly divided fronds. Semi-spreading, attractive ornamental fern.		3"	
Matteuccia struthiopteris Ostrich Fern		48"	M, MW, W	part to full shade	Very attractive with divided, elongate fronds held erect. Unusual fertile fronds emerge from the center in early summer and remain attractive through winter.		3"	
Onoclea sensibilis Sensitive Fern		12-24"	MW, W	part sun to full shade	Unusual fern with broad green fronds and separate brown fertile fronds that are attractive year-round. Good for water		3"	

gardens.

like a bunch of cinnamon sticks.

full sun to

full shade

full sun to

full shade

full sun to

full shade

24-36" M, MW, W

24-36" M, MW, W

30-48" M

Very sun tolerant if always moist as in a water garden. Nice

Unusual fern with fertile portion of leaves in the middle of

central leaves. Extremely showy in the spring and holds up

nicely through the summer. Widely adapted to various soil

Unique fern not even recognized as a fern by many. Excellent

for lakescaping and water gardens. Highly attractive foliage.

semi-erect fronds. Fertile fronds emerge from center and look

3"

3"

3"

Osmunda cinnamomea

Osmunda claytoniana

Cinnamon Fern

Interrupted Fern

Osmunda regalis

Royal Fern

	Flowers	Ht.	Moisture	Light	Description	2" Pot	4" Pot	6" Pot +
Shrubs	•		•		•			
Amelanchier laevis Serviceberry / Juneberry	White April to May	to 12'	D, MD, M, MW	full sun to part shade	One of the earliest flowering shrubs. Stays lower with pruning. Bears bluish purple berries that are edible and attractive to wildlife. Dry to moist soils.			•
Aronia melanocarpa Chokeberry	White May	to 6'	MD, M, MW, W	full to part sun	Very attractive spreading shrub. Nice clusters of white flowers followed by black berries. Deep red fall foliage. Excellent for wet lake edge plantings. Mesic to wet soil.			
Ceanothus americanus New Jersey Tea	White June to Aug	1-2'	D, MD, M	full sun to part shade	Under-utilized, very attractive low shrub with glossy green leaves and 2" clusters of fragrant flowers. Great for foundation plantings and dry restoration sites. Found on dry prairies throughout the region.	•		
Cephalanthus occidentalis Buttonbush	White July to Aug	4-8'	MW, W	full sun to part shade	Unique tropical-looking shrub with glossy foliage and attractive flower clusters in summer. Native range limited to southeastern MN and southwestern WI.			•
Comptonia peregrina Sweet Fern	Brown April to May	1-2'	D, MD, M, MW, W	full sun to part shade	Very cool low shrub with serrated elongate leaves that look like fern leaflets. Very pleasant, fragrant spicy foliage had a number of traditional medicinal uses. One of the most underrated native shrubs.			
Cornus alternifolia Pagoda Dogwood	White May to June	to 16'	MD, M, MW	full sun to part shade	Very attractive shrub with arching leaves that give the appearance of a giant bonsai. Perfect for the Japanese garden as well as habitat restoration or foundation plantings. Wide ranging in MN and WI.			•
Cornus racemosa Gray Dogwood	White May to July	6-8'	MD, M	full sun to part shade	Widely adapted shrub that forms large clonal stands of scattered stems. Good fall color, excellent production of berries. Shade-tolerant.			•
Cornus sericea (stolonifera) Red Osier Dogwood	White May to Aug	to 6'	M, MW, W	full sun to part shade	Very common in wetlands along lakes, streams, and rivers, particularly in the north. Red stems are beautiful in the winter. Mesic to wet soils.			
Corylus americanus Hazelnut	Purple March to April	6-8'	, ,	full sun to part shade	Growing in popularity as landscape plant due to attractive foliage and fall color. Tolerates dry to wet soils and shade to sun. Very good, fast grower.			•
<i>Diervilla lonicera</i> Dwarf Bush Honeysuckle	Yellow June to Sept	2-3.5'	D, MD	full sun to shade	Fantastic low shrub for mass plantings and woodland and forest restorations. Green-bronze arrow-shaped leaves and yellow trumpet-shaped flowers through much of the summer. Dry to moist soils throughout the region.	•	•	•
Hammemalis virginiana Witch Hazel	Yellow October	15'	M, MW	part to full shade	Fall flowering shrub with attractive foliage. Suitable for mesic to moist forest sites. Highly recommended for woodland restoration.			
Ilex verticillata Winterberry Holly	Greenish June	to 6'	M, MW, W	full sun to part shade	Gorgeous in winter. Female plants bear numerous red berries along stems. Excellent lake edge species, but also tolerates most upland soils. Plant at least four to ensure proper sex			
Rhus typhina Staghorn Sumac	Green-yellow July	to 25'	D, MD, M	full sun to part shade	An aggressively suckering shrub great for naturalizing in wilder areas or can be contained by regular removal of sprouts. Beautifully artistic form and texture if left to mature. Dark red clusters of berries stay on all winter & are great food			
Sambucus canadensis Wild Elderberry	White June to July	6-12'	MD, M, MW	full to part sun	Fast-growing shrub tolerant of mesic to wet soils. White flower clusters are followed by deep purple berries. Very attractive and good in jam and wine.			
Spiraea alba Meadowsweet	White June to Aug	3-4'	MW, W	full to part sun	Shrub that frequents lake margins on soil ranging from sand to peat and saturated to moderately dry. Very fast growing for habitat restoration. Great butterfly plant.			•
Spiraea tomentosa Steeplebush	Pink July to Aug	2.5- 3.5'	MW, W	full to part sun	Shrub of lake margins ranging from sandy to peaty. Beautiful tall spikes of pink flowers above attractive foliage. Fast growing for shoreland restoration. Good butterfly plant.			
Symphoricarpos albus Snowberry	White June	3-4'	MD, M	full sun to part shade	Excellent compact shrub for understory plantings and restoration sites. Blue green foliage and abundant white berries are attractive. Found throughout region.			•
Vaccinium angustifolium Lowbush Blueberry	White May to June	6-18"	MD, M, MW	full sun to shade	Yummy, fruit-producing, sprawling plants that grow in a range of habitats ranging from dry, sandy pine barrens to saturated peatlands. Wide-ranging in MN and WI.			•
Viburnum dentatum Arrowwood Viburnum	White June	10-12'	M, MW, W	full sun to part shade	Glossy green toothed leaves. Dark purple berries in summer. Nice tall shrub for shoreland planting or as a hedge. Mesic to wet soils.			
Viburnum lentago Nannyberry	White June	10-15'	M, MW, W	full sun to part shade	Tall shrub with excellent purple fall foliage. Blue-black berries are very tasty fresh or in jam. Great bird cover and forage. Mesic to wet soils.			

	Flowers	ш	Moisture	Light	Description	2" Pot	4" Pot	6" Pot +
Viburnum trilobum Highbush Cranberry	White June	<b>Ht.</b> 10-12'	M, MW, W	full sun to part shade	Poscription Forms tall dense clusters with maple-like lobed leaves. Broad clusters of flowers followed by red berries that persist through winter for beauty and birds.	2	4	•
Trees								
Acer rubrum Red Maple	Red April to May	to 50'	M, MW	full sun to part shade	Mid- to late successional tree, mostly of the northwoods but also into the deciduous forests of central WI and southern MN. Nice fall color, dry to moist soil.			
Acer saccharum Sugar Maple	Yellow April to May	50-75'	M, MW	full sun to part shade	Dominant climax (late successional) tree of decidous forests throughout the region. Best for maple sugaring. Mesic to moist soils.			
Betula nigra River Birch	Brown April to May	60-90'	M, MW, W	full sun to part shade	Cinnamon to reddish peeling bark. Long lived and disease resistant. A very nice tree for shoreland restorations on mesic to wet soils.			
Betula papyrifera Paper Birch	Brown April to May	to 80'	MD, M, MW, W	full sun to part shade	Loved for its white, peeling bark. Gorgeous in clumps on soils ranging from dry to wet. Widely adapted, found throughout the region.			
Carpinus caroliniana Blue Beech/ Musclewood	Green May to June	10-25'	M, MW	full sun to full shade	Tall shrub to small tree with unique rippled wood that resembles striated muscle. Branches tend to go up and out horizontally like an umbrella. Very nice in the landscape.			
Celtis occidentalis Hackberry	Green May	to 60'	MD, M, MW, W	full sun to part shade	Highly adaptable to a variety of soil conditions and urban pollution, this tree is being used more and more as a street/boulevard tree. Very unique corky bark and an open upright spreading form similar to elms.			
Larix laricina Tamarack (American Larch)	none	40-60'	MW, W	full sun to part shade	The only deciduous native conifer. Soft, light-green needles turn a beautiful gold in fall. One of last to lose leaves.  Excellent for lake edge planting as well as in the landscape.  Tolerates all but the driest soils.			
Ostrya virginiana Ironwood/ Hophornbeam	Yellow May to June	15-50'	MD, M, MW	full sun to shade	Slender small tree of sugar maple/basswood forests. Attractive platey bark and very hard wood. Inconspicuous flowers give rise to a hop-like cluster of fruits that persist into			
Picea glauca White Spruce	none	50-75'	М	full sun to part shade	Climax species of the northern deciduous/conifer forests into the northern boreal forests. Very attractive tree for restoration and landscaping.			
Pinus strobus Eastern White Pine	none	100- 150'	MD, M, MW	full sun to part shade	Fast-growing, long-lived pine which brought the first wave of European loggers to the area. Beautiful in the landscape.  Tolerant to soils from wet to dry.			
Populus tremuloides Quaking Aspen	Yellow- Green May	To 50'	MD,M,MW	full to part sun	One of the fastest growing native trees in northern WI it can put on up to 2' of growth per year. Whitish-grey bark adds interest to the landscape.			
Quercus alba White Oak	Green June	to 75'	MD, M, MW	full sun to part shade	Ornamental hardwood valued for its timber and acorn production. Common throughout the region on moist to dry soils.			
Quercus bicolor Swamp White Oak	Yellow- Green May	50-60'	MD, M, MW, W	full sun	More tolerant of wet soils than any other oak, it also adapts well as a boulevard or shade tree in your average yard. Very similar to White Oak.			
Quercus borealis Northern Red Oak	Green June	to 80'	D, MD, M, MW	full sun to part shade	Rapidly growing ornamental hardwood. Best oak for beautiful red fall color. Tolerates moist to dry soils, long-lived.			
Quercus ellipsoidalis Northern Pin Oak	Green June	to 75'	D, MD, M, MW	full to part sun	Pioneer species growing following stand removal through fire or logging. Attractive deeply divided leaves and small acorns in abundance. Particularly adapted to dry soils.			
Quercus macrocarpa Bur Oak	Green June	70-80'	D, MD, M, MW	full sun to part shade	Climax tree species in drier forested areas across the region. Stately, spreading trees common in savannas. Acorns with distinct fringed cap.			
Sorbus decora Showy Mountain Ash Thuja occidentalis	White June to July none	10-30' to 50'	MD, M, MW M, MW, W	full to part sun full sun to	Abundant bright red berries develop in late summer and persist into winter. Berries are loved by birds.  Tree from which many ornamental Arbor Vitae shrubs are			·
Eastern White Cedar				part shade	selected. Forms an attractive pyramid-shaped tree. Good evergreen privacy screen. Valuable wildlife cover.			·
Tilia americana American Linden	Yellow June	to 80'	MD, M, MW	full to part sun	A handsome ornamental shade tree commonly found growing alongside maples in Wiconsin's northern deciduous forests.			•