

DRAGONFLY GARDENS

www.dragonflygardens.net 715-554-7219 natives@dragonflygardens.net Part Sun/Part Shade Medium to Wet Soil Woodland Edge

	Flowers Ht.			AVAILABILITY, SIZE and PRICE subject to	2" Pot	Pot	Pot +	
		Ht.	lt. Moisture	Light	Description		4"	99
Wildflowers							•	
Acorus calamus	Yellow	24-36"	MW, W	full sun to	Unusual plant with attractive yellowish foliage and sweet			т —
Sweet Flag	July to Aug	24-30	10100, 00	part shade	fragrance when crushed. Forms beautiful clumps on lake margins. Great for water gardens.	•		
Anemone canadensis Canada Anemone	White May to July	12-24"	M, MW	full sun to part shade	Fine, showy clump-forming plant of mesic to moist prairies, forest margins and woodlands.	•		
Aster lateriflorus Calico Aster	White Aug to Oct	24"	MD, M, MW	Full sun to shade	Good for short-height restorations in a variety of soil and light conditions. It is an abundant late-season food source for	•	•	
Aster puniceus	Purple	36-56"	MW, W	full sun to	bees. Common, showy aster of lake edges, swamps, bogs, and			
Swamp Aster	Aug to Oct	<u> </u>		part shade	fens. Occurs throughout MN and WI.			$oldsymbol{ol}}}}}}}}}}}}}}}}}}$
Caltha palustris	Yellow	12-18"	MW, W	part sun to	Harbinger of spring. Usually associated with peaty ground			
Marsh Marigold	May to June			shade	and some flowing water, but can be planted on any wet ground. Great for pond edges.		3"	
Campanula americana	Blue	36-60"	MD, M, MW	part sun to	A biennial so grows quickly and flowers the first or second			
Tall Bellflower	July to Oct	00 00	1012, 101, 10100	part shade	year and then self-seeds. Tall spires of violet-blue flowers good for quick color in partially shaded restorations and a variety of soil conditions except yery dry	•		
Chelone glabra	White	24-36"	MW. W	full sun to	Spikes of showy turtle-shaped flowers atop glossy dark green			
Turtlehead	July to Oct		,	part shade	foliage. Nice for gardens and lake edge restorations.	•		
Eupatorium maculatum	Purple	48-60"	MW. W	full sun to	Widespread in MN and WI. Good lake edge perennial with broad flat-topped clusters of			H
Spotted Joe-Pye Weed	July to Sept	40-00		part shade	purple heads. Excellent butterfly plant. Ranges throughout MN and WI	•	•	
Eupatorium perfoliatum	White	36-48"	MW, W	full sun to	Nice lake edge plants with flat-topped cluster of white heads.			
Boneset	June to Aug		,	part shade	Very nice purplish foliage with opposite fused leaves. Used medicinally to hasten healing of bones. Ranges throughout MN and WI	•		
Fragaria vesca	White	6-10"	MD, M, MW	full sun to	Attractive groundcover that grows rapidly. A must for the			
Woodland Strawberry	May to June		, ,	shade	woodland restoration. Sweet tasting red fruits in June are a bonus. Ranges throughout the world in northern latitudes		3"	
Fragaria virginiana	White	6"	D, MD, M,	full sun to	including all of MN and WI Stoloniferous spreading groundcover bears delicious fruit.			H
Wild Strawberry	May to June		MW	shade	Found throughout MN and WI ranging from prairies to forests.	•	•	
Calium baraala	White	16-24"	MD, M,	full sun to	Widely adapted to dry, moist or even wet soils.			-
Galium boreale Northern Bedstraw	June to July	10-24	MW. W	full shade	Common understory species of northern forests in MN & WI. Easy to grow on mesic to dry soils.	•	•	ĺ
Helenium autumnale	Yellow	36-48"	MW, W	full sun to	Sneezeweed won't make you sneeze - it was used as an			
Sneezeweed	July to Oct			shade	expectorant by Native Americans. Gaillardia -like heads in abundance. Good butterfly plant.	•		
Heliopsis helianthoides	Yellow	48-60"	MD, M, MW	full to	Widespread on dry to wet soils. Long flowering period.			
False/Early Sunflower	June to Oct			part sun	Suited to all but very shady restoration sites. Commonly used as a garden perennial.	•	•	
Iris versicolor	Blue	24-36"	M, MW, W	full to	Common on lake margins throughout northern WI and MN.			Г
Northern Blue Flag Iris	June to July		,,	part sun	Very attractive, forming large clumps. Some consider this to	•	•	
Labalia aardinalia	Dod	26 40"	NA)A/ \A/	full ours to	be synonymous with <i>Iris virginica</i> .			\vdash
<i>Lobelia cardinalis</i> Cardinal Flower	Red July to Oct	30-48°	MW, W	full sun to part shade	Showy spikes of red flowers are the best for attracting hummingbirds. Suited to moist soils particularly along	•	•	
Lobelia siphilitica	Blue	20.30"	M, MW, W	full sun to	waterways. Native through most of WI. Widely adapted to wet and moist soils. Vivid blue and white-			\vdash
Great Blue Lobelia	July to Oct	20-30	IVI, IVIVV, VV	part shade	striped flowers in terminal spikes. Compact and well suited to be perennial gardens.	•	•	
Mertensia virginiana	Blue	18-24"	M, MW	part to	Clusters of bell-shaped blue flowers in spring. Nice woodland			Г
Virginia Bluebells	May to June			full shade	wildflower. Dies back by midsummer, but are plantable while		3"	ĺ
=					dormant.			1

Mimulus ringens Monkeyflower	Lavender July to Sept	18-24"	MW, W	full sun to part shade	Lavender spikes of 1" snapdragon flowers that look like small monkeys. Compact plants. Excellent for lakescaping.	•	•	
Physostegia virginiana Obedient Plant	Pink July to Oct	24-36"	M, MW, W	full sun to part shade	Popular garden perennial native to moist prairies, lakeside and streamside wetlands.			
Rudbeckia subtomentosa Sweet Black Eyed	Yellow July to Sept	24-36"	MD, M, MW	full to part sun	Very attractive yellow heads in profusion atop deep green glossy leaves. Good for restorations and perennial gardens. Native into SW WI.	•	•	•
Susan Rudbeckia triloba Brown Eyed Susan	Yellow July to Oct	36-60"	MD, M, MW	full sun to part shade	Outstanding coneflower with numerous heads which are smaller than other species, but more abundant. Very nice foliage. Should be grown more widely in perennial beds. Mesic to wet soils	•	•	
Smilacina racemosa False Solomon Seal	White May to June	18-24"	MD, M, MW	full to part shade	Two-foot tall leafy stemmed lily with a terminal cluster of small white flowers, each of which becomes a red berry. Found throughout forest supporting regions of MN and WI.		3"	
Trillium grandiflorum Trillium	White May	8-12"	M, MW	part shade to shade	Spectacular 2" white 3-petaled flowers in May. This harbinger of spring will let you know that winter is finally over! Widespread in WI and MN forests.		3"	
Verbena hastata Blue Vervain	Blue June to Sept		M, MW, W	full sun to part shade	Tall spikes of blue flowers really attract butterflies. Great for shoreland restorations on mesic to saturated soils. Throughout MN and WI			
Vernonia fasciculata Ironweed	Rose-Pink July to Sept	48-60"	M, MW	full sun to part shade	Broad, flat clusters of pink heads to 5" across. Very attractive to butterflies. Good in moist to wet soils for lake edge			
Veronicastrum virginianum	White June to Sept		MD, M, MW	full sun to part shade	landscaping. Wide ranging in MN and WI. Several tall showy spikes on top of each stem make this an excellent tall garden perennial. Attracts butterflies. Good in		•	
Culver's Root Zizia aurea Golden Alexanders	Yellow May to July	18-30"	MD, M, MW	full sun to part shade	most soils except externely dry. Early-flowering plant of mesic to moist habitats with 3" flat- topped heads of yellow flowers. Seed heads that follow are	•	•	
Grasses, se	daes	and	rushe	<u> </u>	lattractive throughout the year.			
Calamagrostis canadensis	Straw July to Aug		M, MW, W	full sun to part shade	Fine-leaved grass forms a dense cover on lake margins. One of the grasses displaced by reed canary grass and a good			
Blue Joint Reedgrass Carex bebbii Bebb's Sedge	Green May to June	24-36"	M, MW	full sun to part shade	one to plant after nuking RCG. Purplish foliage. Forms dense clumps on moist to saturated soil. Very common in most wetland areas throughout MN and WI.	•		
Carex comosa Bottlebrush Sedge	Green May to June	24-42"	MW, W	full sun to part shade	Beautiful clumps of yellowish stems on lake margins. Nodding flower spikes look like a bottlebrush. Excellent for restorations and water gardens.	•		
Carex crinita Caterpillar Sedge	Green May to June	30-42"	MW, W	full sun to part shade	Clump-forming. 4" nodding spikes resemble a caterpillar. Very nice for wet site restorations and for water gardens. Excellent for control of shoreland erosion.	•		
Carex gracillima Graceful Sedge Carex hystricina	Green May to June Green		MW, W	full sun to full shade full sun to	Clump forming plant of mesic to moist forests throughout MN and WI. Attractive nodding flower spikes. Very similar to Carex comosa. Very attractive plants for	•	٠	
Porcupine Sedge	May to June		,	part shade	shoreland restorations and water gardening.	•		
Carex lupulina Hop Sedge	Green May to June	24-42"	MW, W	full sun to part shade	Flower spikes resemble hops clusters. Stout-stemmed clump former of a wide variety of wet habitats, including shady swamps. Very good for shady rain gardens.	•		
Carex stipata Awl Fruit Sedge	Green May to June	16-40"	MW, W	full sun to full shade	Common, coarse clump forming sedge found in a wide variety of wet habitats throughout MN and WI. Very good fast grower. Shade tolerant species.			
Carex stricta Tussock Sedge	Green May to June	24-36"	MW, W	full sun to part shade	Forms dense clumps of closely clustered fine stems in a variety of wetlands and lake margins. Good shoreland and water garden plant.	•		
Carex vulpinoidea Fox Sedge	Green May to June	18-30"	MD, M, MW, W	full sun to part shade	Beautiful clump-forming sedge of lake and wetland margins. Very attractive multi-tiered flower spikes persist through the summer and into the fall.		•	
Glyceria striata Fowl Manna Grass	Green July to Aug	24-36"	M, MW, W	full sun to full shade	Low clump forming grass common next to small pools in swamps, in moist to seasonally wet forests, and along waterways.	•		
Juncus effusus Soft Rush	Brown July to Aug	24-48"	M, MW, W	full sun to part shade	Deep green wirey stems in large clumps along waterways and in a variety of wetlands. Evergreen plants remain attractive into winter.	•		
Juncus tenuis Path Rush	Brown June to July	to 16"	D, MD, M, MW	full sun to part shade	Very common, widespread plant of wet places throughout MN and WI. Nice for restoration site margins. Easy to grow. Forms attractive clumps.	•		
Scirpus atrovirens Green Bulrush	Brown June to July	36-48"	MW, W	full sun to part shade	Coarse clump forming plant with nice clusters of small brown heads. Suitable for wet shoreland zones.	•		
l	1	1		1				

Scirpus cyperinus	Brown	36-48"	MW, W	full sun to	Coarse clump former with brown fuzzy heads that appear		- 1	
Wool Grass	June to July	30-46	IVIVV, VV	part shade	wooly. One of the most common lake edge species			
Woor Oraco	ourio to oury			partoriado	throughout the region.			
Ferns								
Onoclea sensibilis		12-24"	MW, W	part sun to	Unusual fern with broad green fronds and separate brown		1	
Sensitive Fern			,	full shade	fertile fronds that are attractive year-round. Good for water		3"	
					gardens.			
Osmunda cinnamomea		24-36"	M, MW, W	full sun to	Very sun tolerant if always moist as in a water garden. Nice		2"	
Cinnamon Fern				full shade	semi-erect fronds. Fertile fronds emerge from center and look like a bunch of cinnamon sticks.		3"	
Osmunda regalis		24-36"	M, MW, W	full sun to	Unique fern not even recognized as a fern by many. Excellent			
Royal Fern			, ,	full shade	for lakescaping and water gardens. Highly attractive foliage.		3"	
Shrubs								
Amelanchier laevis	White	to 12'	D, MD, M,	full sun to	One of the earliest flowering shrubs. Stays lower with pruning.			
Serviceberry /	April to May		MW	part shade	Bears bluish purple berries that are edible and attractive to			•
Juneberry Aronia melanocarpa	White	to 6'	MD, M,	full to part	wildlife. Drv to moist soils. Very attractive spreading shrub. Nice clusters of white flowers			
Chokeberry	May	10 0	MW, W	sun	followed by black berries. Deep red fall foliage. Excellent for			
one we serif	····ay		·		wet lake edge plantings. Mesic to wet soil.			
Cephalanthus	White	4-8'	MW, W	full sun to	Unique tropical-looking shrub with glossy foliage and			
occidentalis	July to Aug			part shade	attractive flower clusters in summer. Native range limited to			•
Buttonbush Comptonia peregrina	Brown	1-2'	D, MD, M,	full sun to	southeastern MN and southwestern WI. Very cool low shrub with serrated elongate leaves that look			
Sweet Fern	April to May	l -	MW, W	part shade	like fern leaflets. Very pleasant, fragrant spicy foliage had a			
	'		,	ľ	number of traditional medicinal uses. One of the most under-			٠
O-maria	\A/I- :4 -	4- 01	NA NAVA/ VA/	full access to	rated native shrubs			
Cornus sericea (stolonifera)	White May to Aug	to 6'	M, MW, W	full sun to part shade	Very common in wetlands along lakes, streams, and rivers, particularly in the north. Red stems are beautiful in the winter.			
Red Osier Dogwood	May to Aug			part snaue	Mesic to wet soils.			
Hammemalis virginiana	Yellow	15'	M, MW	part to full	Fall flowering shrub with attractive foliage. Suitable for mesic			
Witch Hazel	October			shade	to moist forest sites. Highly recommended for woodland			•
llex verticillata	Greenish	to 6'	M, MW, W	full sun to	restoration. Gorgeous in winter. Female plants bear numerous red berries			
Winterberry Holly	June	10 6	IVI, IVIVV, VV	part shade	along stems. Excellent lake edge species, but also tolerates			
Transcriborry Fromy	Guno			partoriado	most upland soils. Plant at least four to ensure proper sex			•
					ratio			
Sambucus canadensis	White	6-12'	MD, M, MW	full to part	Fast-growing shrub tolerant of mesic to wet soils. White			
Wild Elderberry	June to July			sun	flower clusters are followed by deep purple berries. Very attractive and good in iam and wine.			·
Vaccinium	White	6-18"	MD, M, MW	full sun to	Yummy, fruit-producing, sprawling plants that grow in a range			
angustifolium	May to June			shade	of habitats ranging from dry, sandy pine barrens to saturated			•
Lowbush Blueberry Viburnum dentatum	White	10 10'	M, MW, W	full sun to	peatlands. Wide-ranging in MN and WI. Glossy green toothed leaves. Dark purple berries in summer.			
Arrowwood Viburnum	June	10-12	IVI, IVIVV, VV	part shade	Nice tall shrub for shoreland planting or as a hedge. Mesic to			
7 (ITOWWOOd VIDalilaiii	dune			part snade	wet soils.			
Viburnum lentago	White	10-15'	M, MW, W	full sun to	Tall shrub with excellent purple fall foliage. Blue-black berries			
Nannyberry	June			part shade	are very tasty fresh or in jam. Great bird cover and forage.			•
Viburnum trilobum	White	10-12'	M, MW, W	full sun to	Mesic to wet soils. Forms tall dense clusters with maple-like lobed leaves. Broad			
Highbush Cranberry	June	10 12	,,	part shade	clusters of flowers followed by red berries that persist through			•
,					winter for beauty and birds.			
Trees								
Betula nigra	Brown	60-90'	M, MW, W	full sun to	Cinnamon to reddish peeling bark. Long lived and disease			
River Birch	April to May		,,	part shade	resistant. A very nice tree for shoreland restorations on mesic			•
					to wet soils.			
Betula papyrifera	Brown	to 80'	MD, M,	full sun to	Loved for its white, peeling bark. Gorgeous in clumps on soils			_
Paper Birch	April to May		MW, W	part shade	ranging from dry to wet. Widely adapted, found throughout the region.			•
Larix laricina	none	40-60'	MW, W	full sun to	The only deciduous native conifer. Soft, light-green needles			
Tamarack				part shade	turn a beautiful gold in fall. One of last to lose leaves.			
(American Larch)					Excellent for lake edge planting as well as in the landscape.			
Ostrya virginiana	Yellow	15-50'	MD, M, MW	full sun to	Tolerates all but the driest soils Slender small tree of sugar maple/basswood forests.	\vdash		
Ironwood/	May to June	.5 50	, 171, 17177	shade	Attractive platey bark and very hard wood. Inconspicuous			_
Hophornbeam	.,				flowers give rise to a hop-like cluster of fruits that persist into			•
·		FC ==:	L	f	the winter	\sqcup		
Picea glauca	none	50-75'	IVI	full sun to	Climax species of the northern deciduous/conifer forests into			
White Spruce				part shade	the northern boreal forests. Very attractive tree for restoration and landscaping.			-
Thuja occidentalis	none	to 50'	M, MW, W	full sun to	Tree from which many ornamental Arbor Vitae shrubs are			
Eastern White Cedar				part shade	selected. Forms an attractive pyramid-shaped tree. Good			•
			l	<u> </u>	evergreen privacy screen. Valuable wildlife cover.			