

## DRAGONFLY GARDENS

Native Catalog 2022 Edition

www.dragonflygardens.net 715-554-7219 natives@dragonflygardens.net

491 Hwy 46 Amery, WI

AVAILABILITY, SIZE and PRICE subject to change.

|  | Flowers                      | Ht.         | Moisture        | Light                     | Description  | 2" Pot | 4" Pot                |
|--|------------------------------|-------------|-----------------|---------------------------|--|--------|-----------------------|
| Wildflowers  |                              | 1           |                 | <u> </u>                  | 1-000p.1101.   |        |                       |
| Achillea millefolium<br>Yarrow                       | White<br>June to Oct         | 18-30"      | D, MD, M,<br>MW | full sun to<br>part shade | Attractive, aromatic, ferny foliage. Great component of a no-mow or low-mow lawn. Good butterfly plant. Grows well in a wide variety of habitats.  |        |                       |
| Acorus calamus<br>Sweet Flag                         | Yellow<br>July to Aug        | 24-<br>36"  | MW, W           | full sun to<br>part shade | Unusual plant with attractive yellowish foliage and sweet fragrance when crushed. Forms beautiful clumps on lake margins. Great for water gardens.   | •      |                       |
| <i>Agastache foeniculum</i><br>Anise Hyssop          | Blue<br>June to Oct          | 24-40"      | MD, M           | full sun to<br>part shade | Beautiful 3" flower spikes attract butterflies throughout summer.<br>Excellent cutflower and nice in perennial gardens. Fragrant foliage<br>makes good herbal tea.   | •      |                       |
| Allium cernuum<br>Nodding Wild Onion                 | Pink<br>July to Sept         | 12-18"      | MD, M, MW       | full to<br>part sun       | Nodding 2" flower clusters above thin, wispy stems. Edible flower bulbs.   | •      | $\lceil \cdot \rceil$ |
| Allium stellatum<br>Prairie Onion                    | Pink<br>July to Sept         | 12-18"      | D, MD, M,<br>MW | full to<br>part sun       | Attractive plants occurring in dry prairies and on rocky cliff areas.  Good plant for tough sites and rock gardens. Ranging throughout  MN and WI.   |        |                       |
| Amorpha canescens<br>Leadplant                       | Blue<br>June to July         | 18-42"      | D, MD, M        | full sun to<br>part shade | Long-lived shrub with attractive fine silvery foliage. Slow-growing but worth the wait. Used medicinally by Native Americans. Suited to driest sandy soil.   |        |                       |
| Anaphilis margaritacea Pearly Everlasting            | White<br>June to Aug         | 18-24"      | D, MD           | full sun to<br>part shade | White papery strawflowers produced atop attractive whitish foliage.  Nice dried flower and contrast plant in the perennial garden. Host plant for painted lady butterfly.                                    |        |                       |
| Anemone canadensis<br>Canada Anemone                 | White<br>May to July         | 12-24"      | M, MW           | full sun to<br>part shade | Fine, showy clump-forming plant of mesic to moist prairies, forest margins and woodlands.  | •      |                       |
| Anemone cylindrica<br>Thimbleweed                    | White<br>Late May to<br>July | 24-<br>36"  | D, MD, M        | full sun to<br>part shade | Thimble-like infloresence held atop slender stems above highly divided basal leaves. Very pretty from summer into fall. Good on very dry sandy soils.  |        |                       |
| Antennaria<br>plantaginifolia<br>Plantain Pussy Toes | White<br>May to July         | 6-12"       | D, DM, M        | full to part<br>sun       | Nice groundcover for dry habitats in all but full shade. Attractive white foliage, excellent lawn alternative for tough sites. Ranges through MN and WI.   |        |                       |
| Aquilegia canadensis<br>Wild Columbine               | Orange<br>May to June        | 20-36"      | D, MD, M        | full sun to<br>shade      | Free-flowering woodland perennial that forms large clumps with attractive foliage. Excellent hummingbird plant. Often referred to as "honeysuckle."  |        | •                     |
| <i>Artemesia ludoviciana</i><br>Prairie Sage         | Yellow<br>July to Oct        | 24-<br>48"  | D, MD, M        | full to<br>part sun       | Fragrant whitish foliage makes a nice contrast to green plants. Dry to mesic prairies. Sacred incense and insect repellant of Native Americans.  |        |                       |
| <i>Asarum canadense</i><br>Wild Ginger               | Maroon<br>May to June        | 6-12"       | MD, M, MW       | part sun to<br>full shade | Beautiful forest groundcover with nearly round leaves. Strong ginger fragrance. Unusual flowers held on ground said to be pollinated by beetles. A must for forest restorations and shady perennial gardens. |        | 3"                    |
| <i>Asclepias incarnata</i><br>Swamp Milkweed         | Lavender<br>June to Aug      | 36-48"      | M, MW, W        | full sun to<br>part shade | Showy plant of moist to wet habitats. Host plant for monarch butterflies, very good nectar source for butterflies and hummingbirds. Great for lake edges.  |        | •                     |
| <i>Asclepias syriaca</i><br>Common Milkweed          | Pink<br>June to Aug          | 36"-<br>60" | D, MD, M,<br>MW | full sun to<br>part sun   | Large leaved, robust plant known for being the larval host for Monarch butterflies. Benefits many other pollinators as well. Easy and fast to establish.   |        |                       |
| <i>Asclepias tuberosa</i><br>Butterfly Weed          | Orange<br>June to Sept       | 12-24"      | D, MD, M        | full to<br>part sun       | A must for gardens, producing orange flowers in 3" clusters. Tolerates driest sandy soils. Host plant for monarch butterflies and nectar source for others.  |        |                       |
| Asclepias verticillata<br>Whorled Milkweed           | White<br>July to Sept        | 12-24"      | D, MD, M        | full to part<br>sun       | This delicate and fine-leaved plant is rather inconspicuous in the landscape but like other milkweeds it serves as an important food source for Monarch caterpillars. Tolerant of dry soils.                 |        |                       |

|  | Flowers                 | Ht.    | Moisture         | Light                     | Description  | 2" Pot | 4" Pot |
|--|-------------------------|--------|------------------|---------------------------|--|--------|--------|
| Aster laevis   | Blue                    |        | MD, M, MW        | full sun to               | Very showy vibrant blue-flowered perennial. Several cultivated   | 2      | 4      |
| Smooth Aster   | July to Oct             |        | IVID, IVI, IVIVV | part shade                | varieties exist, attesting to its beauty. Great butterfly plant and cutflower.   | •      |        |
| Aster lateriflorus<br>Calico Aster                         | White<br>Aug to Oct     | 24"    | MD, M, MW        | Full sun to shade         | Good for short-height restorations in a variety of soil and light conditions. It is an abundant late-season food source for bees.  |        |        |
| Aster macrophyllus Big Leaf Aster                          | Blue<br>Aug to Oct      | 8-18"  | D, MD, M,<br>MW  | full sun to<br>shade      | One of the most common groundcovers of northern forests. Broad heart-shaped leaves make an great, fast-growing groundcover for all but the wettest soil types.   |        |        |
| Aster novae-angliae<br>New England Aster                   | Blue<br>Aug to Oct      | 36-60" | MD, M, MW,<br>W  | full sun to<br>part shade | Nice perennial for moist to wet habitats. Commonly used as a garden perennial and for wet lake-edge plantings. Nectar source for butterflies.  |        | •      |
| Aster oblongifolius<br>Aromatic Aster                      | Violet<br>Aug to Oct    | 12-24" | D, MD            | full to<br>part sun       | Very attractive low-grower for dry sandy prairies. Common throughout SE to NW MN and SW to NW WI. A must for short prairie restorations.   | •      |        |
| Aster oolentangiense<br>Sky Blue Aster                     | Blue<br>Aug to Oct      |        | D, MD, M         | full to part<br>sun       | An aster with one of the truest blue flowers you can find among native wildflowers. Attracts loads of pollinators.   |        | •      |
| Aster ptarmicoides<br>Upland White Aster                   | White<br>Aug to Oct     | 12-24" | D, MD            | full to<br>part sun       | Low, spreading plants covered with white heads late summer through<br>fall. Mesic to dry prairies throughout MN and WI. Excellent in short<br>prairies.  |        |        |
| Aster puniceus<br>Swamp Aster                              | Purple<br>Aug to Oct    | 36-56" | MW, W            | full sun to<br>part shade | Common, showy aster of lake edges, swamps, bogs, and fens. Occurs throughout MN and WI.  | •      |        |
| Aster umbellatus<br>Flat-topped Aster                      | White<br>July to Aug    | 30-42" | M, MW, W         | full sun to<br>full shade | White heads held in a broad terminal cluster. Common plant of moist to wet lake edges and moist to mesic forests. Great butterfly plant and cutflower. Ranges throughout MN and WI.  |        |        |
| <i>Baptisia alba</i><br>White Wild Indigo                  | White<br>June to Aug    | 30-42" | D, MD, M,<br>MW  | full to<br>part sun       | Legume of moist to wet prairies and lake edges. Attractive foliage and flowers followed by pods. Nice in perennial gardens.  |        |        |
| Baptisia australis<br>Blue False Indigo                    | Blue<br>June to July    | 30-42" | M, MW            | full to<br>part sun       | Stately perennial with glossy foliage and beautiful sky blue flower clusters. Excellent garden perennial. Native south of our range but tough enough for restorations here in NW WI.   |        |        |
| Callirhoe triangulata<br>Clustered Poppy Mallow            | Pink<br>July to Sept    | 12"    | D, MD            | full to part<br>sun       | Thrives in dry, gravely soil and produces exceptionally beautiful magenta-red blossoms. Also known as "Wine Cups."   |        |        |
| <i>Caltha palustris</i><br>Marsh Marigold                  | Yellow<br>May to June   | 12-18" | MW, W            | part sun to<br>shade      | Harbinger of spring. Usually associated with peaty ground and some flowing water, but can be planted on any wet ground. Great for pond edges.  |        | 3"     |
| Campanula americana<br>Tall Bellflower                     | Blue<br>July to Oct     | 36-60" | MD, M, MW        | part sun to<br>part shade | A biennial so grows quickly and flowers the first or second year and then self-seeds. Tall spires of violet-blue flowers good for quick color in partially shaded restorations and a variety of soil conditions except very dry. |        |        |
| Campanula rotundifolia<br>Harebell                         | Blue<br>June to Sept    | 16-24" | D, MD            | full sun to<br>part shade | Blue bell-shaped flowers atop delicate, grassy foliage. Great plant for restorations. Common throughout WI in all dry habitats. Shade-tolerant.  |        | 3"     |
| Chelone glabra<br>Turtlehead                               | White<br>July to Oct    |        | MW, W            | full sun to<br>part shade | Spikes of showy turtle-shaped flowers atop glossy dark green foliage. Nice for gardens and lake edge restorations. Widespread in MN and WI.  | •      |        |
| Coreopsis lanceolata Lance Leaf Coreopsis                  | Yellow<br>June to Aug   |        | D, MD            | full sun to part shade    | Wiry stems bearing yellow daisies. Easy to grow on toughest, driest soil types. Very good butterfly plant.   | •      | •      |
| Coreopsis palmata<br>Prairie Coreopsis                     | Yellow<br>June to Oct   | 16-30" | D, MD, M         | full sun to<br>part shade | Yellow daisies above ferny foliage. Native of driest prairies and bluff communities throughout southern 2/3 of MN and WI. Good for butterflies.  |        |        |
| <i>Dalea candida</i><br>White Prairie Clover               | White<br>June to Sept   | 16-30" | D, MD, M         | full to<br>part sun       | Dense 2" clusters of small white flowers atop grayish ferny foliage.<br>Very attractive perennial for prairie and savanna restorations as well<br>as perennial gardens.  |        |        |
| Dalea purpurea<br>Purple Prairie Clover                    | Purple<br>June to Sept  | 16-30" | D, MD, M         | full to<br>part sun       | Similar to <i>Dalea candida</i> but with purple flowers. Both species are excellent butterfly plants. Used as tea by Native Americans.   |        |        |
| Echinacea angustifolia<br>Narrow-Leaf Purple<br>Coneflower | Lavender<br>July to Aug | 24-36" | D, MD            | full to<br>part sun       | Said to be the best of the purple coneflowers for medicinal purposes. Native in Western MN and further west, but hardy in this area. Very attractive with narrow leaves.   |        | •      |
| Echinacea pallida<br>Pale Purple Coneflower                | Lavender<br>July to Aug | 36-48" | D, MD, M         | full to<br>part sun       | Only echinacea species definitely native to WI. Nice narrow foliage.<br>Heads have reflexed rays ("petals"). Can't beat these for butterflies.   | •      |        |
| Echinacea purpurea Purple Coneflower                       | Purple<br>July to Aug   | 36-48" | MD, M, MW        | full to<br>part sun       | Spectacular display of broad-headed purple daisies. Great for restorations and perennial gardens.  |        | •      |

|  | Eleviere                 | L14    | Majatura              | l iada4                   | Decarintian  | " Pot       | " Pot       |
|--|--------------------------|--------|-----------------------|---------------------------|--|-------------|-------------|
| Eryngium yuccifolium                                     | Flowers<br>White         | Ht.    | Moisture<br>MD, M, MW | Light<br>full to          | Description     Exotic-looking plant that resembles a desert yucca. Bristly white  | 2           | 4           |
| Rattlesnake Master                                       | July to Oct              |        |                       | part sun                  | heads and whitish foliage make it a great background perennial as well as a tough fast grower for restorations.  | ·           | ·           |
| Eupatorium maculatum<br>Spotted Joe-Pye Weed             | Purple<br>July to Sept   | 48-60" | MW, W                 | full sun to part shade    | Good lake edge perennial with broad flat-topped clusters of purple heads. Excellent butterfly plant. Ranges throughout MN and WI.  |             |             |
| Eupatorium perfoliatum<br>Boneset                        | White<br>June to Aug     | 36-48" | MW, W                 | full sun to<br>part shade | Nice lake edge plants with flat-topped cluster of white heads. Very nice purplish foliage with opposite fused leaves. Used medicinally to hasten healing of bones. Ranges throughout MN and WI.                        | •           |             |
| Eupatorium purpureum<br>Purple Joe Pye Weed              | Purple<br>July to Sept   | 48-72" | MD, M, MW             | part sun to<br>shade      | Moist to moderately dry savannas and forests through southern 2/3 of WI and eastern 1/2 of MN. Attractive broad clusters of purple heads. Excellent for attracting butterflies.  |             |             |
| Euthamia graminfolia<br>Grass-leaved Goldenrod           | Yellow<br>July to Sept   | 30-48" | MD, M, MW,<br>W       | Full sun                  | This plant is not really a goldenrod but is an excellent source of pollen for bees. Can be somewhat spreading and aggressive but is very adaptable to a variety of soil moisture conditions and good for restorations. |             |             |
| Fragaria virginiana<br>Wild Strawberry                   | White<br>May to June     | 6"     | D, MD, M,<br>MW       | full sun to<br>shade      | Stoloniferous spreading groundcover bears delicious fruit. Found throughout MN and WI ranging from prairies to forests. Widely adapted to dry, moist or even wet soils.  |             |             |
| Galium boreale<br>Northern Bedstraw                      | White<br>June to July    |        | MD, M, MW,<br>W       | full shade                | Common understory species of northern forests in MN & WI. Easy to grow on mesic to dry soils.  | ·           | •           |
| Geranium maculatum<br>Wild Geranium                      | Pink<br>May to June      |        | MD, M                 | full to<br>part shade     | Tolerant to all soils and woodland conditions except deeply shaded situations. Nice perennial geranium with attractive foliage that turns red in fall.   |             | 3"          |
| Geum triflorum<br>Prairie Smoke                          | Red<br>May to June       | 6-12"  | D, MD, M,<br>MW       | full to<br>part sun       | Awesome early-flowering prairie wildflower, great for sandy site restorations and for rock gardening. Hairy fruits remain attractive throughout the year.  |             |             |
| Helenium autumnale<br>Sneezeweed                         | Yellow<br>July to Oct    | 36-48" | MW, W                 | full sun to<br>shade      | Sneezeweed won't make you sneeze - it was used as an expectorant by Native Americans. Gaillardia -like heads in abundance. Good butterfly plant.   |             |             |
| Heliopsis helianthoides<br>False/Early Sunflower         | Yellow<br>June to Oct    | 48-60" | MD, M, MW             | full to<br>part sun       | Widespread on dry to wet soils. Long flowering period. Suited to all but very shady restoration sites. Commonly used as a garden perennial.  | •           | •           |
| Heuchera richardsonii<br>Alum Root                       | White<br>May to July     | 12-18" | D, MD, M,<br>MW       | full sun to<br>part shade | Common through most of WI and MN. Attractive, tough ground cover well suited to dry soils. Quite shade-tolerant. Usually associated with rock outcrops in prairies, woodlands, and forests.                            |             |             |
| Iris versicolor<br>Northern Blue Flag Iris               | Blue<br>June to July     | 24-36" | M, MW, W              | full to<br>part sun       | Common on lake margins throughout northern WI and MN. Very attractive, forming large clumps. Some consider this to be synonymous with <i>Iris virginica</i> .  |             |             |
| <i>Liatris aspera</i><br>Rough Blazing Star              | Purple<br>July to Sept   | 18-36" | D, MD, M              | full to<br>part sun       | Beautiful purple spikes. Most common liatris of dry, sandy prairies throughout MN and WI. Excellent butterfly plant.   | •           | •           |
| Liatris ligulistylis<br>Northern Plains Blazing<br>Star  | Purple<br>July to Aug    | 36-60" | MD, M, MW             | full to<br>part sun       | Rosy heads appear in abundance on tall spikes. Beautiful plant among the best to attract butterflies. OK on wet and dry soils.   |             |             |
| <i>Liatris pychnostachya</i><br>Thick Spike Blazing Star | Purple<br>July to Sept   | 36-60" | M, MW, W              | full to<br>part sun       | Dense purple spikes. Good for moist and wet restoration sites. Very showy, good butterfly plant. Ranges throughout prairie regions of MN and WI.   |             |             |
| <i>Liatris spicata</i><br>Marsh Blazing Star             | Purple<br>July to Aug    | 36-60" | M, MW, W              | full to<br>part sun       | Most common liatris for perennial gardening. Grows well in any condition. Perfect pollinator plant. Native into SE WI. Moist to wet soils.   |             |             |
| Lobelia cardinalis<br>Cardinal Flower                    | Red<br>July to Oct       | 36-48" | MW, W                 | full sun to<br>part shade | Showy spikes of red flowers are the best for attracting hummingbirds. Suited to moist soils particularly along waterways. Native through most of WI.   |             |             |
| Lobelia siphilitica<br>Great Blue Lobelia                | Blue<br>July to Oct      | 20-30" | M, MW, W              | full sun to<br>part shade | Widely adapted to wet and moist soils. Vivid blue and white-striped flowers in terminal spikes. Compact and well suited to perennial gardens.  |             |             |
| <i>Mertensia virginiana</i><br>Virginia Bluebells        | Blue<br>May to June      | 18-24" | M, MW                 | part to<br>full shade     | Clusters of bell-shaped blue flowers in spring. Nice woodland wildflower. Dies back by midsummer, but are plantable while dormant.   |             | 3"          |
| Mimulus ringens<br>Monkeyflower                          | Lavender<br>July to Sept |        | MW, W                 | full sun to<br>part shade | Lavender spikes of 1" snapdragon flowers that look like small monkeys. Compact plants. Excellent for lakescaping.  | •           | •           |
| <i>Monarda fistulosa</i><br>Bergamot                     | Lavender<br>July to Aug  | 36-48" | D, MD, M,<br>MW       | full sun to<br>part shade | One of the most common and widespread native wildflowers. Good in herbal teas. Used medicinally by Native Americans. Moist to dry soils.   | $  \cdot  $ | $  \cdot  $ |

|  | Flowers                          | Ht.    | Moisture        | Light                                 | Description   | 2" Pot | 4" Pot |
|--|----------------------------------|--------|-----------------|---------------------------------------|---|--------|--------|
| Monarda punctata<br>Dotted Mint                        | Lavender<br>June to Sept         |        | D, MD           | full to<br>part sun                   | Fragrant herb with showy dotted multicolored leaves below flowers. Good on dry soils. Used medicinally and in herbal teas. Nice cutflower.  |        |        |
| <i>Oenothera pilosella</i><br>Prairie Sundrops         | Yellow<br>May to June            | 24"    | MD, M, MW       | full to part<br>sun                   | A day-flowering species of the Evening Primrose family. It spreads semi-agressively but can easily be kept under control due to shallow roots. Flowers attract pollinators.                                     |        |        |
| Penstemon digitalis<br>Smooth Penstemon                | White<br>June to July            | 30-48" | MD, M           | full sun to<br>part shade             | 1.5" white tubular flowers in tall spikes. Good garden perennial and cutflower. Tolerant of dry and mesic soils and a variety of light conditions.  |        |        |
| Phlox divaricata<br>Woodland Phlox                     | Blue<br>June to July             | 12-18" | MD, M, MW       | part sun to<br>shade                  | Beautiful fragrant spring wildflower occurring in forests ranging from mesic to floodplain. Excellent for shady restorations and gardens.   |        |        |
| Phlox pilosa   | Pink                             | 18-24" | D, MD, M,       | full to                               | Broad heads of deep pink flowers. Very nice wildflower for  |        |        |
| Prairie Phlox Physostegia virginiana Obedient Plant    | May to June Pink July to Oct     | 24-36" | MW<br>M, MW, W  | part sun<br>full sun to<br>part shade | restorations and perennial gardens.  Popular garden perennial native to moist prairies, lakeside and streamside wetlands.   |        |        |
| Polemonium reptans<br>Jacob's Ladder                   | Blue<br>May to June              | 18"    | MD, M, MW       | part sun to<br>full shade             | Beautiful spring woodland wildflower. Ferny leaves resemble a ladder. A must for woodland gardens and restorations.   |        |        |
| Ratibida pinnata<br>Yellow Coneflower                  | Yellow<br>July to Oct            | 24-52" | MD, M           | full to<br>part sun                   | One of the more common native prairie wildflowers. Very attractive to butterflies. Greyish foliage makes this a nice contrast plant.  |        |        |
| Rudbeckia fulgida<br>Orange Coneflower                 | Orange-<br>Yellow<br>July to Oct | 24-36" | MD, M, MW       | full to<br>part sun                   | Beautiful compact coneflower with deep green foliage and 3" orange-<br>yellow heads with brown centers. Parent of 'goldsturm', one of the<br>most popular garden perennials.                                    |        |        |
| <i>Rudbeckia hirta</i><br>Black Eyed Susan             | Yellow<br>June to Oct            | 24-36" | MD, M, MW       | full to<br>part sun                   | Very common short-lived perennial occurring in a wide range of habitats. Produces lots of color the first year.   | •      |        |
| Rudbeckia laciniata Wild Goldenglow                    | Yellow<br>July to Sept           | 48-60" | M, MW           | full sun to<br>part shade             | Found throughout WI, primarily in swamp habitats and along lakes and streams, but also in prairies. Good for moist, shaded restoration sites.   |        |        |
| Rudbeckia<br>subtomentosa<br>Sweet Black Eyed<br>Susan | Yellow<br>July to Sept           | 24-36" | MD, M, MW       | full to<br>part sun                   | Very attractive yellow heads in profusion atop deep green glossy leaves. Good for restorations and perennial gardens. Native into SW WI.  |        |        |
| Rudbeckia triloba<br>Brown Eyed Susan                  | Yellow<br>July to Oct            | 36-60" | MD, M, MW       | full sun to<br>part shade             | Outstanding coneflower with numerous heads which are smaller than other species, but more abundant. Very nice foliage. Should be grown more widely in perennial beds. Mesic to wet soils.                       |        |        |
| Silene regia<br>Royal Catchfly                         | Red<br>June to Oct               | 18-24" | MD, M           | full sun to<br>part shade             | Deep red tubular flowers flaring out into a star. Excellent hummingbird plant and garden perennial. Native just south of MN and WI, but hardy in the north.   |        | •      |
| Silphium perfoliatum<br>Cup Plant                      | Yellow<br>July to Sept           | 48-72" | M, MW           | full sun to<br>part shade             | Impressive sunflower with numerous broad yellow heads. Unusual opposite leaves fuse to make cups which collect water for bird bathing. Birds love it!   |        |        |
| Solidago flexicaulis<br>Zig Zag Goldenrod              | Yellow<br>Aug to Sept            | 24"    | MD, M, MW       | part to<br>full shade                 | Good woodland perennial for restorations. Unique zig-zag stems with scattered yellow heads. Very different than most goldenrods. Moist to mesic soils.  |        |        |
| Solidago nemoralis<br>Gray Goldenrod                   | Yellow<br>Aug to Oct             | 24-36" | D, MD           | Sun to part<br>shade                  | One of the most attractive goldenrods, it has vibrant golden-yellow flowers that are a great source of pollen for bees and butterflies right up to frost. It is also one of the most shade-tolerant goldenrods. |        |        |
| Solidago rigida<br>Stiff Goldenrod                     | Yellow<br>Aug to Sept            | 36-48" | D, MD, M,<br>MW | full to<br>part sun                   | Another goldenrod worthy of growing for its very attractive foliage. Topped with a broad, flat cluster of yellow heads. Adapted to mesic and moist soils. Good garden perennial.                                |        |        |
| Solidago speciosa<br>Showy Goldenrod                   | Yellow<br>Aug to Sept            | 24-48" | D, MD, M        | full sun to<br>part shade             | Highly recommended for restorations and for perennial gardens. Very showy tall heads above glossy green foliage. Great butterfly plant.   |        |        |
| <i>Symphyotrichum shortii</i><br>Short's Aster         | Blue<br>Aug to Oct               | 36"    | MD, M           | full to part<br>shade                 | And easy way to get a profusion of warm-blue flowers in your shade garden or restoration. Similar in use and habitat to Big Leaf Aster, looks great with Zig Zag Goldenrod.                                     |        |        |
| <i>Tradescantia ohiensis</i><br>Ohio Spiderwort        | Blue<br>June to Oct              | 24-48" | D, MD, M,<br>MW | full sun to<br>part shade             | Forms large clumps creating a very nice display of 1" flowers in heads on top of each stem. Great for restorations and gardens. Moist to dry soils.   |        |        |
| <i>Verbena hastata</i><br>Blue Vervain                 | Blue<br>June to Sept             | 48-60" | M, MW, W        | full sun to<br>part shade             | Tall spikes of blue flowers really attract butterflies. Great for shoreland restorations on mesic to saturated soils. Throughout MN and WI.   |        |        |
| <i>Verbena stricta</i><br>Hoary Vervain                | Blue<br>June to Sept             | 24-36" | D, MD           | full sun to<br>part shade             | Spikes of blue flowers above greyish foliage. Adapted to dry, sandy soils. Good butterfly plant.  |        | Ŀ      |
|  |                                  |        |                 |                                       |   |        |        |

|   | Flowers                   | Ht.    | Moisture  | Light                     | Description  | 2" Pot | 4" Pot   |
|---|---------------------------|--------|-----------|---------------------------|--|--------|----------|
| Vernonia fasciculata<br>Ironweed              | Rose-Pink<br>July to Sept |        |           | full sun to<br>part shade | Broad, flat clusters of pink heads to 5" across. Very attractive to butterflies. Good in moist to wet soils for lake edge landscaping. Wide ranging in MN and WI.                  |        | <u> </u> |
| Veronicastrum<br>virginianum<br>Culver's Root | White<br>June to Sept     |        | MD, M, MW | full sun to<br>part shade | Several tall showy spikes on top of each stem make this an excellent tall garden perennial. Attracts butterflies. Good in most soils except externely dry.                         |        |          |
| <i>Viola pedatifida</i><br>Prairie Violet     | Blue<br>May to June       | 4"     | MD, M     | full sun to<br>part shade | 1.5" blue flowers above deeply clawed foliage. Very nice for edging restoration sites on dry to mesic soils. Also nice for rock gardening. Common in prairie regions of MN and WI. |        | 3"       |
| <i>Zizia aurea</i><br>Golden Alexanders       | Yellow<br>May to July     | 18-30" | MD, M, MW | full sun to<br>part shade | Early-flowering plant of mesic to moist habitats with 3" flat-topped heads of yellow flowers. Seed heads that follow are attractive throughout the year.                           | •      |          |

|   |                                   |             |                 |                           |   | Pot | Pot |
|---|-----------------------------------|-------------|-----------------|---------------------------|---|-----|-----|
|   | Flowers                           | Ht.         | Moisture        | Light                     | Description   | 5   | 4   |
| Grasses, se   | edges,                            | and         | rushe           | S                         |   |     |     |
| Andropogon gerardii<br>Big Bluestem                 | Purplish<br>Green<br>July to Sept | 36-<br>72+" | D, MD, M,<br>MW | full sun to<br>part shade | One of the most dominant tall prairie grasses. Purplish foliage creates a nice effect along with turkey-foot-like flower head. Dry to moist habitats.   |     |     |
| Bouteloua curtipendula<br>Side Oats Grama           | Red-Green<br>July to Sept         | 18-30"      | D, MD, M        | full sun to<br>part shade | Very common species of dry to mesic habitats. Stamens are red, making them beautiful cutflowers when fresh. A must for any open dry site restoration.   |     |     |
| <i>Bouteloua gracilis</i><br>Blue Grama             | Purplish<br>Green<br>July to Sept | 12"         | D, MD           | full to<br>part sun       | Native through the central Great Plains into western WI. Very good lawn alternative. Low growing, forming attractive clumps.  |     |     |
| Calamagrostis<br>canadensis<br>Blue Joint Reedgrass | Straw<br>July to Aug              | 36-48"      | M, MW, W        | full sun to<br>part shade | Fine-leaved grass forms a dense cover on lake margins. One of the grasses displaced by reed canary grass and a good one to plant after nuking RCG. Purplish foliage.                                      |     |     |
| Carex bebbii<br>Bebb's Sedge                        | Green<br>May to June              | 24-36"      | M, MW           | full sun to<br>part shade | Forms dense clumps on moist to saturated soil. Very common in most wetland areas throughout MN and WI.  |     |     |
| Carex bicknellii<br>Bicknell's Sedge                | Tan<br>May to June                | 12-24"      | MD, M, MW       | full to<br>part sun       | Sedge adapted to habitats ranging from dry to moist, usually on rocky, gravelly soil. Forms nice clumps. Found throughout southern 2/3 of MN and WI.  | •   |     |
| Carex comosa<br>Bottlebrush Sedge                   | Green<br>May to June              | 24-42"      | MW, W           | full sun to<br>part shade | Beautiful clumps of yellowish stems on lake margins. Nodding flower spikes look like a bottlebrush. Excellent for restorations and water gardens.   | •   |     |
| Carex crinita<br>Caterpillar Sedge                  | Green<br>May to June              | 30-42"      | MW, W           | full sun to<br>part shade | Clump-forming. 4" nodding spikes resemble a caterpillar. Very nice for wet site restorations and for water gardens. Excellent for control of shoreland erosion.   |     |     |
| Carex gracillima<br>Graceful Sedge                  | Green<br>May to June              | 18-30"      | MW, W           | full sun to<br>full shade | Clump forming plant of mesic to moist forests throughout MN and WI. Attractive nodding flower spikes.   | •   |     |
| Carex hystricina Porcupine Sedge                    | Green<br>May to June              |             | MW, W           | full sun to<br>part shade | Very similar to <i>Carex comosa</i> . Very attractive plants for shoreland restorations and water gardening.  | •   |     |
| Carex lupulina<br>Hop Sedge                         | Green<br>May to June              | 24-42"      | MW, W           | full sun to<br>part shade | Flower spikes resemble hops clusters. Stout-stemmed clump former of a wide variety of wet habitats, including shady swamps. Very good for shady rain gardens.   | •   |     |
| Carex pennsylvanica<br>Pennsylvania Sedge           | Brown<br>May                      | 6-8"        | D, MD, M        | full sun to<br>full shade | Outstanding, low, clump-forming graminoid with yellow-green foliage. Forms extensive stands in northwoods forests that are extremely attractive. A must for forest restorations in northern MN and WI.    | •   |     |
| Carex radiata<br>Eastern Star Sedge                 | Green<br>June                     | 12"         | M, MW           | full to part<br>shade     | Attractive dark green, fine-textured groundcover for rich, moist woods. Not suitable for dry woodlands  | •   |     |
| <i>Carex stipata</i><br>Awl Fruit Sedge             | Green<br>May to June              | 16-40"      | MW, W           | full sun to<br>full shade | Common, coarse clump forming sedge found in a wide variety of wet habitats throughout MN and WI. Very good fast grower. Shade tolerant species.   |     |     |
| Carex stricta<br>Tussock Sedge                      | Green<br>May to June              | 24-36"      | MW, W           | full sun to<br>part shade | Forms dense clumps of closely clustered fine stems in a variety of wetlands and lake margins. Good shoreland and water garden plant.  |     |     |
| Carex vulpinoidea<br>Fox Sedge                      | Green<br>May to June              | 18-30"      | MD, M, MW,<br>W | full sun to<br>part shade | Beautiful clump-forming sedge of lake and wetland margins. Very attractive multi-tiered flower spikes persist through the summer and into the fall.   |     | ·   |
| Elymus hystrix (patula)<br>Bottlebrush Grass        | Green<br>June to July             | 36-48"      | MD, M           | full sun to<br>full shade | Clump forming grass with tall spikes topped with an infloresence that looks like a bottlebrush. A must for forest restorations and shady perennial gardens.   |     |     |
| <i>Glyceria striata</i><br>Fowl Manna Grass         | Green<br>July to Aug              | 24-36"      | M, MW, W        | full sun to<br>full shade | Low clump forming grass common next to small pools in swamps, in moist to seasonally wet forests, and along waterways.  |     |     |
| Hierochloe odorata<br>Sweet Grass                   | Green<br>May to June              | 12-24"      | M, MW, W        | full sun to<br>part shade | Widespread in MN and WI mostly on wetland margins on moist to wet soils. Sacred plant for Native Americans, used for braiding and basket-weaving. Revered for its sweet, aromatic vanilla-like fragrance. |     |     |
| Juncus effusus<br>Soft Rush                         | Brown<br>July to Aug              | 24-48"      | M, MW, W        | full sun to<br>part shade | Deep green wirey stems in large clumps along waterways and in a variety of wetlands. Evergreen plants remain attractive into winter.  |     |     |
| <i>Juncus tenuis</i><br>Path Rush                   | Brown<br>June to July             | to 16"      | D, MD, M,<br>MW | full sun to<br>part shade | Very common, widespread plant of wet places throughout MN and WI. Nice for restoration site margins. Easy to grow. Forms attractive clumps.   |     |     |
| Koeleria macrantha<br>June Grass                    | Green<br>June to July             | 12-24"      | D, MD           | full sun to<br>part shade | Clump-forming grass, most commonly associated with dry sandy soils. Low-grower suitable for edging native restoration plantings.  |     | ·   |

|   | Flowers               | Ht.    | Moisture        | Light                     | Description   | 2" Pot | 4" Pot |
|---|-----------------------|--------|-----------------|---------------------------|---|--------|--------|
| Panicum virgatum<br>Switch Grass              | Purple<br>June to Oct | 36-48" | D, MD, M,<br>MW | full to<br>part sun       | Fast growing clump former with attractive open infloresence. Suited to soils ranging from mesic to wet. Good for steep banks and slopes.                              |        |        |
| Schizachyrium<br>scoparium<br>Little Bluestem | Amber<br>July to Sept | 18-36" | D, MD, M        | full sun to<br>part shade | Fabulous ornamental grass native to dry, sandy to mesic habitats throughout the region. Amber foliage maintains color through winter. Forms neat clumps.              | •      |        |
| Scirpus atrovirens<br>Green Bulrush           | Brown<br>June to July | 36-48" | MW, W           | full sun to<br>part shade | Coarse clump forming plant with nice clusters of small brown heads. Suitable for wet shoreland zones.   |        |        |
| Scirpus cyperinus<br>Wool Grass               | Brown<br>June to July | 36-48" | MW, W           | full sun to<br>part shade | Coarse clump former with brown fuzzy heads that appear wooly. One of the most common lake edge species throughout the region.   |        |        |
| Sorghastrum nutans<br>Indian Grass            | Amber<br>July to Sept | 48-72" | D, MD, M        | full sun to<br>part shade | Clump-forming grass of dry, sandy to moist habitats. Very ornamental and suitable for the back of perennial borders. Great plant for a wide range of habitats.        |        |        |
| Spartina pectinata<br>Prairie Cordgrass       | Yellow<br>Aug to Oct  | 48-72" | M, MW, W        | full to<br>part sun       | Rhizomatous grass forming large clumps in sandy soil, usually near the water table, but sometimes under dry conditions. Common shoreland plant in northern WI and MN. |        |        |
| Sporobolus heterolepis<br>Prairie Dropseed    | Green<br>Aug to Oct   | 18-36" | D, MD, M,<br>MW | full to<br>part sun       | Clump former with open, fine sprays of flowers. Popular as a garden perennial. Native to dry and mesic prairies throughout the region.                                |        |        |

## Ferns

|                       |        |           |              |   | <br> |
|-----------------------|--------|-----------|--------------|---|------|
| Adiantum pedatum      | 24"    | MD, M, MW | I.           | Graceful stems with soft highly divided leaves in a U-shaped pattern. |      |
| Maidenhair Fern       |        |           | shade        | Slow-starting but eventually forms nice clumps. Moist to wet forests. | 3"   |
| Athyrium felix-femina | 36"    | M, MW     | part to full | Semi-spreading lime green highly divided leaves make this a very      | 3"   |
| Lady Fern             |        |           | shade        | useful ground cover for shady restorations and gardens.               | ١٦   |
| Dryopteris marginalis | 20"    | MD, M, MW | part to full | Blue-green highly divided fronds. Semi-spreading, attractive          | 3"   |
| Eastern Wood Fern     |        |           | shade        | ornamental fern.  | 3    |
| Matteuccia            | 48"    | M, MW, W  | part to full | Very attractive with divided, elongate fronds held erect. Unusual     |      |
| struthiopteris        |        |           | shade        | fertile fronds emerge from the center in early summer and remain      | 3"   |
| Ostrich Fern          |        |           |              | attractive through winter.  |      |
| Onoclea sensibilis    | 12-24" | MW, W     | part sun to  | Unusual fern with broad green fronds and separate brown fertile       |      |
| Sensitive Fern        |        |           | full shade   | fronds that are attractive year-round. Good for water gardens.        | 3"   |
| Osmunda cinnamomea    | 24-36" | M, MW, W  | full sun to  | Very sun tolerant if always moist as in a water garden. Nice semi-    |      |
| Cinnamon Fern         |        |           | full shade   | erect fronds. Fertile fronds emerge from center and look like a bunch | 3"   |
|                       |        |           |              | of cinnamon sticks.   |      |

|  | Flowers                     | Ht.          | Moisture           | Light                     | Description   | 1 Gallon       | 2 Gallon | 3 Gallon |
|--|-----------------------------|--------------|--------------------|---------------------------|---|----------------|----------|----------|
| Shrubs   |                             |              |                    |                           |   |                |          |          |
| Amelanchier laevis<br>Serviceberry / Juneberry         |                             | to 12'       | D, MD, M,<br>MW    | full sun to<br>part shade | One of the earliest flowering shrubs. Stays lower with pruning. Bears bluish purple berries that are edible and attractive to wildlife. Dry to moist soils.   | •              |          |          |
| <i>Aronia melanocarpa</i><br>Chokeberry                | White<br>May                | to 6'        | MD, M, MW,<br>W    | full to part<br>sun       | Very attractive spreading shrub. Nice clusters of white flowers followed by black berries. Deep red fall foliage. Excellent for wet lake edge plantings. Mesic to wet soil.   |                |          |          |
| Ceanothus americanus<br>New Jersey Tea                 | White<br>June to Aug        | 1-2'         | D, MD, M           | full sun to<br>part shade | Under-utilized, very attractive low shrub with glossy green leaves and 2" clusters of fragrant flowers. Great for foundation plantings and dry restoration sites. Found on dry prairies throughout the region.                                      | 2"<br>4"       |          |          |
| Cephalanthus<br>occidentalis<br>Buttonbush             | White<br>July to Aug        | 4-8'         | MW, W              | full sun to<br>part shade | Unique tropical-looking shrub with glossy foliage and attractive flower clusters in summer. Native range limited to southeastern MN and southwestern WI.  |                | •        |          |
| Comptonia peregrina<br>Sweet Fern                      | Brown<br>April to May       | 1-2'         | D, MD, M,<br>MW, W | full sun to<br>part shade | Very cool low shrub with serrated elongate leaves that look like fern leaflets. Very pleasant, fragrant spicy foliage had a number of traditional medicinal uses. One of the most underrated native shrubs.   | •              |          |          |
| Cornus alternifolia<br>Pagoda Dogwood                  | White<br>May to June        | to 16'       | MD, M, MW          | full sun to<br>part shade | Very attractive shrub with arching leaves that give the appearance of a giant bonsai. Perfect for the Japanese garden as well as habitat restoration or foundation plantings. Wide ranging in MN and WI.  | •              |          |          |
| Cornus racemosa<br>Gray Dogwood                        | White<br>May to July        | 6-8'         | MD, M              | full sun to<br>part shade | Widely adapted shrub that forms large clonal stands of scattered stems. Good fall color, excellent production of berries. Shade-tolerant.   | •              | •        |          |
| Cornus sericea<br>(stolonifera)<br>Red Osier Dogwood   | White<br>May to Aug         | to 6'        | M, MW, W           | full sun to<br>part shade | Very common in wetlands along lakes, streams, and rivers, particularly in the north. Red stems are beautiful in the winter. Mesic to wet soils.   | •              |          |          |
| Corylus americanus<br>Hazelnut                         | Purple<br>March to<br>April | 6-8'         | MD, M, MW          | part shade                | Growing in popularity as landscape plant due to attractive foliage and fall color. Tolerates dry to wet soils and shade to sun. Very good, fast grower.   | •              |          |          |
| <i>Diervilla Ionicera</i><br>Dwarf Bush<br>Honeysuckle | Yellow<br>June to Sept      | 2-3.5'       | D, MD              | full sun to<br>shade      | Fantastic low shrub for mass plantings and woodland and forest restorations. Green-bronze arrow-shaped leaves and yellow trumpet-shaped flowers through much of the summer. Dry to moist soils throughout the region.                               | 2"<br>4"<br>1G |          |          |
| <i>Hammemalis virginiana</i><br>Witch Hazel            | Yellow<br>October           | 15'          | M, MW              | part to full<br>shade     | Fall flowering shrub with attractive foliage. Suitable for mesic to moist forest sites. Highly recommended for woodland restoration.  |                |          |          |
| <i>llex verticillata</i><br>Winterberry Holly          | Greenish<br>June            | to 6'        | M, MW, W           | full sun to<br>part shade | Gorgeous in winter. Female plants bear numerous red berries along stems. Excellent lake edge species, but also tolerates most upland soils. Plant at least four to ensure proper sex  |                |          |          |
| Rhus typhina<br>Staghorn Sumac                         | Green-yellow<br>July        | to 25'       | D, MD, M           | full sun to<br>part shade | An aggressively suckering shrub great for naturalizing in wilder areas or can be contained by regular removal of sprouts. Beautifully artistic form and texture if left to mature. Dark red clusters of berries stay on all winter & are great food |                |          |          |
| Sambucus canadensis<br>Wild Elderberry                 | White<br>June to July       | 6-12'        |                    | full to part<br>sun       | Fast-growing shrub tolerant of mesic to wet soils. White flower clusters are followed by deep purple berries. Very attractive and good in jam and wine.   | •              |          |          |
| <i>Spiraea alba</i><br>Meadowsweet                     | White<br>June to Aug        | 3-4'         | MW, W              | full to part<br>sun       | Shrub that frequents lake margins on soil ranging from sand to peat and saturated to moderately dry. Very fast growing for habitat restoration. Great butterfly plant.  |                |          |          |
| <i>Spiraea tomentosa</i><br>Steeplebush                | Pink<br>July to Aug         | 2.5-<br>3.5' | MW, W              | full to part<br>sun       | Shrub of lake margins ranging from sandy to peaty. Beautiful tall spikes of pink flowers above attractive foliage. Fast growing for shoreland restoration. Good butterfly plant.  |                |          |          |
| Symphoricarpos albus<br>Snowberry                      | White<br>June               | 3-4'         | MD, M              | full sun to<br>part shade | Excellent compact shrub for understory plantings and restoration sites. Blue green foliage and abundant white berries are attractive. Found throughout region.  | •              | •        |          |
| Vaccinium<br>angustifolium<br>Lowbush Blueberry        | White<br>May to June        | 6-18"        | MD, M, MW          | full sun to<br>shade      | Yummy, fruit-producing, sprawling plants that grow in a range of habitats ranging from dry, sandy pine barrens to saturated peatlands. Wide-ranging in MN and WI.   | •              |          |          |
| Viburnum dentatum<br>Arrowwood Viburnum                | White<br>June               | 10-12'       | M, MW, W           | full sun to<br>part shade | Glossy green toothed leaves. Dark purple berries in summer. Nice tall shrub for shoreland planting or as a hedge. Mesic to wet soils.   |                |          |          |
| <i>Viburnum lentago</i><br>Nannyberry                  | White<br>June               | 10-15'       | M, MW, W           | full sun to<br>part shade | Tall shrub with excellent purple fall foliage. Blue-black berries are very tasty fresh or in jam. Great bird cover and forage. Mesic to wet soils.  |                | •        | •        |

|   | Flamera                 |                   | Majotumo             | l imbé                    | Description   | Gallon | Gallon | Gallon |
|---|-------------------------|-------------------|----------------------|---------------------------|---|--------|--------|--------|
| Viburnum trilobum<br>Highbush Cranberry           | Flowers White June      | <b>Ht.</b> 10-12' | Moisture<br>M, MW, W | full sun to<br>part shade | Poscription  Forms tall dense clusters with maple-like lobed leaves. Broad clusters of flowers followed by red berries that persist through winter for beauty and birds.  | 1      | 2      |        |
| Trees   | •                       | •                 |                      |                           | + Indicates larger potted trees us  | sually | avai   | lable  |
| Acer rubrum<br>Red Maple                          | Red<br>April to May     | to 50'            | M, MW                | full sun to<br>part shade | Mid- to late successional tree, mostly of the northwoods but also into the deciduous forests of central WI and southern MN. Nice fall color, dry to moist soil.   |        |        | +      |
| <i>Acer saccharum</i><br>Sugar Maple              | Yellow<br>April to May  |                   | M, MW                | full sun to<br>part shade | Dominant climax (late successional) tree of decidous forests throughout the region. Best for maple sugaring. Mesic to moist soils.  | •      |        | +      |
| Betula nigra<br>River Birch                       | Brown<br>April to May   | 60-90'            | M, MW, W             | full sun to<br>part shade | Cinnamon to reddish peeling bark. Long lived and disease resistant. A very nice tree for shoreland restorations on mesic to wet soils.  |        | •      | +      |
| <i>Betula papyrifera</i><br>Paper Birch           | Brown<br>April to May   | to 80'            | MD, M, MW,<br>W      | full sun to<br>part shade | Loved for its white, peeling bark. Gorgeous in clumps on soils ranging from dry to wet. Widely adapted, found throughout the region.  |        | •      | +      |
| Carpinus caroliniana<br>Blue Beech/<br>Musclewood | Green<br>May to June    | 10-25'            | M, MW                | full sun to<br>full shade | Tall shrub to small tree with unique rippled wood that resembles striated muscle. Branches tend to go up and out horizontally like an umbrella. Very nice in the landscape.   | •      |        | +      |
| Celtis occidentalis<br>Hackberry                  | Green<br>May            | to 60'            | MD, M, MW,<br>W      | full sun to<br>part shade | Highly adaptable to a variety of soil conditions and urban pollution, this tree is being used more and more as a street/boulevard tree. Very unique corky bark and an open upright spreading form similar to elms.          |        |        |        |
| Larix laricina<br>Tamarack<br>(American Larch)    | none                    | 40-60'            | MW, W                | full sun to<br>part shade | The only deciduous native conifer. Soft, light-green needles turn a beautiful gold in fall. One of last to lose leaves.  Excellent for lake edge planting as well as in the landscape.  Tolerates all but the driest soils. | •      | •      |        |
| Ostrya virginiana<br>Ironwood/<br>Hophornbeam     | Yellow<br>May to June   | 15-50'            | MD, M, MW            | full sun to<br>shade      | Slender small tree of sugar maple/basswood forests. Attractive platey bark and very hard wood. Inconspicuous flowers give rise to a hop-like cluster of fruits that persist into  | •      |        |        |
| Picea glauca<br>White Spruce                      | none                    | 50-75'            | М                    | full sun to<br>part shade | Climax species of the northern deciduous/conifer forests into<br>the northern boreal forests. Very attractive tree for restoration<br>and landscaping.  |        |        |        |
| Pinus strobus<br>Eastern White Pine               | none                    | 100-<br>150'      | MD, M, MW            | full sun to<br>part shade | Fast-growing, long-lived pine which brought the first wave of European loggers to the area. Beautiful in the landscape.  Tolerant to soils from wet to dry.   |        |        | +      |
| Populus tremuloides<br>Quaking Aspen              | Yellow-<br>Green<br>May | To 50'            | MD,M,MW              | full to part<br>sun       | One of the fastest growing native trees in northern WI it can put on up to 2' of growth per year. Whitish-grey bark adds interest to the landscape.   |        |        | +      |
| Quercus alba<br>White Oak                         | Green<br>June           | to 75'            | MD, M, MW            | full sun to<br>part shade | Ornamental hardwood valued for its timber and acorn production. Common throughout the region on moist to dry soils.   |        |        | +      |
| Quercus bicolor<br>Swamp White Oak                | Yellow-<br>Green<br>Mav | 50-60'            | MD, M, MW,<br>W      |                           | More tolerant of wet soils than any other oak, it also adapts well as a boulevard or shade tree in your average yard. Very similar to White Oak.  |        |        | +      |
| Quercus borealis<br>Northern Red Oak              | Green<br>June           | to 80'            | D, MD, M,<br>MW      | full sun to<br>part shade | Rapidly growing ornamental hardwood. Best oak for beautiful red fall color. Tolerates moist to dry soils, long-lived.   |        |        | +      |
| Quercus ellipsoidalis<br>Northern Pin Oak         | Green<br>June           | to 75'            | D, MD, M,<br>MW      | full to part<br>sun       | Pioneer species growing following stand removal through fire or logging. Attractive deeply divided leaves and small acorns in abundance. Particularly adapted to dry soils.   |        |        | +      |
| <i>Quercus macrocarpa</i><br>Bur Oak              | Green<br>June           |                   | D, MD, M,<br>MW      | full sun to<br>part shade | Climax tree species in drier forested areas across the region. Stately, spreading trees common in savannas. Acorns with distinct fringed cap.   |        |        | +      |
| Sorbus decora<br>Showy Mountain Ash               | White<br>June to July   |                   |                      | full to part<br>sun       | Abundant bright red berries develop in late summer and persist into winter. Berries are loved by birds.   |        | •      |        |
| Thuja occidentalis<br>Eastern White Cedar         | none                    | to 50'            | M, MW, W             | full sun to<br>part shade | Tree from which many ornamental Arbor Vitae shrubs are selected. Forms an attractive pyramid-shaped tree. Good evergreen privacy screen. Valuable wildlife cover.   |        | •      |        |
| <i>Tilia americana</i><br>American Linden         | Yellow<br>June          | to 80'            | MD, M, MW            | full to part<br>sun       | A handsome ornamental shade tree commonly found growing alongside maples in Wiconsin's northern deciduous forests.  |        | •      | +      |